


# Endometriosis factsheet



Endorsed by

# What is endometriosis?

Do you see female patients? Do they have painful periods? Pain pre or post their periods? Painful sex? Do they suffer chronic pelvic pain, which may be intermittent or constant? Do they have pain when passing urine or with bowel movements? Is it painful to place a speculum for a smear test?

## Have you considered endometriosis?

One in 10 women have endometriosis and it takes an average of eight years for women to get diagnosed. Endometriosis can often be confused with or misdiagnosed as IBS (irritable bowel syndrome). It is important that women receive an early diagnosis, or working diagnosis, of endometriosis even if hormonal treatments appear to be working, as women need to understand that they have a long-term, chronic condition and the impact this might have on their future and their fertility. The RCN recognises that we all live in a gender diverse society and that this standard may be used by and/or applied to people who identify as non-binary or transgender.

## What is endometriosis?

Endometriosis is defined as the presence of endometrial-like tissue outside the uterus, which induces a chronic, inflammatory reaction, and may lead to scar tissue formation. There are different types of endometriosis including superficial, deep infiltrating disease (>5mm deep into the tissue) and endometrioma.

While some women with endometriosis experience painful symptoms and/or infertility, others have no symptoms at all (or may consider their symptoms to be normal).

The exact prevalence of endometriosis is unknown but the World Health Organization estimates that 10% (WHO, 2021) of the general female population and up to 50% of infertile women have endometriosis.

NICE published guidance in 2017, which details diagnosing and managing endometriosis, and should be consulted for further information.




You can find out more about the condition by taking a look at the informative website [endometriosis-uk.org](https://www.endometriosis-uk.org). Endometriosis UK is a charity that works to improve the lives of people affected by endometriosis and to decrease the impact it has on those with the condition and their families and friends.

## Some quick facts and figures about endometriosis

Facts and figures – from Endometriosis UK  
[endometriosis-uk.org](https://www.endometriosis-uk.org)

- **One in 10** women of reproductive age (between puberty and menopause) in the UK suffer from endometriosis.
- **10%** of women worldwide have endometriosis – that's **190 million worldwide**.
- The prevalence of endometriosis in women with infertility is as high as 50%.
- Endometriosis is the second most common gynaecological condition (after fibroids) in the UK.
- Endometriosis affects **1.5 million women**, a similar number of women are affected by diabetes or asthma.
- On average it takes **8 years** from onset of symptoms to get a diagnosis.
- Endometriosis costs the UK economy **£8.2bn** a year in treatment, loss of education, work and health care costs.
- The cause of endometriosis is unknown, but there are many different treatment options.

Information provided by Endometriosis UK



## Who may be affected?

Women of any age can be affected by endometriosis but it is rare for the condition to manifest before a girl has her first period.

Teenagers who suffer with painful periods, experience fainting or collapse when having a period, or who miss school because of their period problems should be considered as possibly suffering from the condition.

## What are the symptoms?

Symptoms may vary from woman to woman and some women may experience no symptoms at all (or may not recognise their symptoms as abnormal).

Typical endometriosis symptoms include:

- painful periods
- deep pain during sex
- chronic pelvic pain
- painful bowel movements, painful urination and blood in urine
- cyclical or premenstrual symptoms with or without abnormal bleeding and pain. Heavy periods are not a symptom of endometriosis, although a lot of women with endometriosis also have heavy periods.
- chronic fatigue
- depression (depression is not a direct symptom, however may be a side effect of the long diagnosis, or having a chronic condition and infertility).
- a family history of endometriosis
- infertility
- painful caesarean section scar or cyclical lump
- back, legs and chest pain.



Endometriosis should be considered early in young women with pelvic pain as there is often a delay of between seven and 12 years from the onset of symptoms to receiving a definitive diagnosis.



## Care management in primary care

Suspected endometriosis may be managed in primary care but consider referral to gynaecology or a specialist endometriosis centre if there is any suspicion or uncertainty over the cause of pain or if women are presenting with fertility issues. Women with suspected deep endometriosis involving the bowel, bladder or ureter must be referred to a specialist endometriosis service. All women should have as a minimum an abdominal examination and if appropriate a pelvic examination (NICE 2018 quality standard), and should be informed that endometriosis is being considered.

Treatments that can be tried in primary care include:

- **analgesics** – either simple or non-steroidal anti-inflammatory drugs (NSAID); these can be used in combination and especially around the time of the period
- **oral hormonal treatments** – combined oral contraceptive pills can be taken conventionally, continuously without a break, or in a tricycling regimen (three packs together); if women cannot have estrogen then the progesterone-only pill (eg, cerazette) could be used but it is important to remember that not all women will experience amenorrhoea so pain may persist; other alternatives include a course of medroxyprogesterone acetate (MPA), norethisterone or Dienogest (MHRA to approve this drug for use in England is awaited and it is now available to prescribe for endometriosis). If the initial course of hormonal treatment does not manage symptoms, the woman should be referred to a gynaecologist (NICE 2018 quality standard)
- **intra-uterine hormones** – an intra-uterine system such as Mirena IUS may provide relief from pain and is also a long-term treatment.

## When to refer?

If you see a woman with the above symptoms, encourage her to see her GP or consider a referral to gynaecology. Be aware of local arrangements and seek advice from an endometriosis clinical nurse specialist (this may not be not available in all areas, alternatively refer to local gynaecology department):

- if there is uncertainty over the diagnosis
- if a women requests referral



- if the woman has fertility problems
- if surgical and medical management of endometriosis is required
- if complex/severe endometriosis is suspected – for example, endometriomas or where endometriosis is affecting the bowel (NICE 2018 quality standard)
- if initial hormonal treatment for endometriosis is not effective, not tolerated or contraindicated (NICE guidelines 2017 and Quality Standard 2018).

Please note, a six-month timescale can be used to decide whether initial hormonal treatment is effective however a referral should be made before six months if it becomes clear that treatment is not effective.


## Care in secondary care and endometriosis centres

Women with endometriosis often need referral to secondary care for the diagnosis and treatment of the condition.

The investigations offered include ultrasound scan, although a negative scan or MRI does not rule out endometriosis, therefore the gold standard for diagnosis is laparoscopy with biopsy. Laparoscopy can be diagnostic but more often this is combined with operative surgical procedures to remove the endometriosis. Investigations offered should include a specialist ultrasound scan to identify deep infiltrating endometriosis in the pelvis.

Cases of severe endometriosis (or suspected severe endometriosis) should be sent to a specialist BSGE (British Society for Gynaecology Endoscopy) accredited endometriosis centre where women can access specialist gynaecologists and a clinical nurse specialist (CNS) who work in conjunction with general surgeons and urologists. These specialist centres also liaise with pain management teams and have links with a local fertility team.

A full list of accredited specialist endometriosis centres in the UK can be found online at the BSGE website at: [bsge.org.uk](https://www.bsg.org.uk)



## Further information and resources

NICE (2017) *Endometriosis: diagnosis and management NICE guideline* [NG73]. Available at: [nice.org.uk/guidance/ng73](https://www.nice.org.uk/guidance/ng73)

NICE (2018) Endometriosis Quality standard [QS172]. Available at: [nice.org.uk/guidance/qs172](https://www.nice.org.uk/guidance/qs172)

Norton W, Holloway D, Mitchell H and Law C (2020) The role of Endometriosis Clinical Nurse Specialists in British Society for Gynaecological Endoscopy registered centres: A UK survey of practice. *Nursing open*, 7(6), pp.1852-1860. Available at: [ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)

All Party Parliamentary Group on Endometriosis (2020) *Endometriosis in the UK: Time for Change – APPG on Endometriosis Inquiry Report 2020*. Available at: [endometriosis-uk](https://www.endometriosis-uk.org)

To discover more about the condition or access additional information resources, here are some websites together with some topical research studies you might find helpful.

Endometriosis UK [endometriosis-uk.org](https://www.endometriosis-uk.org)

The British Society for Gynaecological Endoscopy [bsge.org.uk](https://www.bsge.org.uk)

RCGP (2021) menstrual wellbeing toolkit – see [rcgp.org.uk](https://www.rcgp.org.uk)

RCN (2020) Promoting menstrual well being – see [rcn.org.uk](https://www.rcn.org.uk)

RCN (2021) Endometriosis – see [rcn.org.uk](https://www.rcn.org.uk)

Royal College of Obstetricians and Gynaecologists [rcog.org.uk](https://www.rcog.org.uk)

European Society of Human Reproduction and Embryology [eshre.eu](https://www.eshre.eu)

WHO (2021) Endometriosis – see [who.int](https://www.who.int)

The World Endometriosis Society [endometriosis.ca](https://www.endometriosis.ca)

The ENDOPART study findings, published in 2013, provide a powerful insight into the emotional, sexual and relational impacts of the disease. See *Endometriosis: improving the wellbeing of couples – summary report and recommendations*. Available at: [dmu.ac.uk/research](https://www.dmu.ac.uk/research)

A 2010 study provides valuable indicators for providing care and services to women from minority ethnic groups. See Endometriosis and cultural diversity: improving services for minority ethnic women at: [cultureandcompassion.com](https://www.cultureandcompassion.com)



It has been suggested that I may have endometriosis.

Symptoms	Yes/no	Notes
Painful periods		
Deep pain during sex		
Chronic pelvic pain (periodic or constant)		
Pain during bowel movements		
Painful urination and/or blood in urine		
Cyclical or premenstrual symptoms with or without abnormal bleeding and pain		
Chronic fatigue		
Depression		
Family history of endometriosis		
Infertility		
Painful caesarean section scar or cyclical lump in the scar		
Pain in back, legs and/or chest		
Heavy periods		Whilst not a direct symptom, many with endometriosis have heavy bleeding



## Endometriosis

Please tear off this section and give it to women who you feel should seek a referral.



The RCN represents nurses and nursing, promotes excellence in practice and shapes health policies.

September 2021

Review date: September 2024

Published by the  
Royal College of Nursing  
20 Cavendish Square  
London W1G 0RN

RCN Direct 0345 772 6100

RCN Online [rcn.org.uk](http://rcn.org.uk)