


Female Genital Mutilation

An RCN resource for nursing and midwifery practice
(Fifth edition)

CLINICAL PROFESSIONAL RESOURCE



Acknowledgements

This fifth edition of *Female Genital Mutilation: An RCN resource for nursing and midwifery practice* was reviewed and updated in 2023 by Ruth Bailey, RCN Womens Health Forum Chair and Carmel Bagness RCN Professional Lead, Midwifery and Womens Health. It builds on numerous contributions from a wide range of health care professionals who successfully contribute to the ever increasing awareness of the impact of FGM, and the need for health care professionals to continue to engage and campaign on behalf of victims and survivors. We thank you all.

The Professional Lead at the RCN Midwifery and Womens Health is Carmel Bagness, carmel.bagness@rcn.org.uk

This document has been designed in collaboration with our members to ensure it meets most accessibility standards. However, if this does not fit your requirements, please contact corporate.communications@rcn.org.uk

RCN Legal Disclaimer

This publication contains information, advice and guidance to help members of the RCN. It is intended for use within the UK but readers are advised that practices may vary in each country and outside the UK. The information in this booklet has been compiled from professional sources, but its accuracy is not guaranteed. Whilst every effort has been made to ensure the RCN provides accurate and expert information and guidance, it is impossible to predict all the circumstances in which it may be used. Accordingly, the RCN shall not be liable to any person or entity with respect to any loss or damage caused or alleged to be caused directly or indirectly by what is contained in or left out of this website information and guidance.

Published by the Royal College of Nursing, 20 Cavendish Square, London, W1G 0RN

© 2023 Royal College of Nursing. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means electronic, mechanical, photocopying, recording or otherwise, without prior permission of the Publishers. This publication may not be lent, resold, hired out or otherwise disposed of by ways of trade in any form of binding or cover other than that in which it is published, without the prior consent of the Publishers.

Contents

Female genital mutilation (FGM)	5
Introduction	5
What is FGM?	7
Historical and cultural context	7
Prevalence	9
Tackling FGM in the UK	11
FGM procedures and health effects	17
Types of FGM	17
Health risks and complications	20
Human rights, legal and professional responsibility	23
A human rights issue	23
Legal aspects	23
Safeguarding now and for future generations	25
Service provision and multi-agency working	28
Communication	28
Multi-agency working	28
Service provision	29
Health visitors, school nurses and community children's nurses	30
Community, practice and travel nurses	31
Acute sector nurses	32
Midwives	32
Nursing associates	33
Confidentiality	33
Referral to appropriate agencies	34
Education, raising awareness and conscious engagement to change	35
Practice matters and procedures	36
Professional curiosity and assessing need	36
Knowledge about diversity	37

Antenatal care and reversal (deinfibulation)	37
Care in labour	39
Conclusion.....	41
Highlighting the campaign to end FGM.....	41
References and further reading.....	42
Resources	46
Organisations and support groups	47
Appendices	49
Appendix 1: Map showing the global distribution of FGM	49
Appendix 2: Alternative terms used for FGM	50
Appendix 3: Overview of mandatory reporting.....	51
Appendix 4: Tackling FGM in England.....	52

Female genital mutilation (FGM)

Introduction

Female genital mutilation (FGM) or cutting, previously referred to as female circumcision, is a challenging subject to understand and manage.

The term female genital mutilation (FGM) refers to all procedures that involve the total or partial removal of the external female genitalia for non medical reasons and is practised in around 31 countries across the world. FGM may result in serious harm, it has no health benefits and is now internationally recognised as a violation of human rights (WHO, 2023).

FGM is usually performed on young girls from infancy to 15 years old, although older women may also be at risk.

It is estimated that around 200 million women and girls alive today have been subjected to FGM with a further three million girls estimated to be at risk of FGM annually.

In 2008, the World Health Assembly passed a resolution to eliminate FGM by 2030, and this remains a global target to protect against violence towards women and girls.

FGM is child abuse and the practice is illegal in the UK. The UN is clear that FGM is torture and calls for its elimination as a form of cruel, inhumane and degrading treatment of girls and women (RCM Position statement last updated 2015). The hidden nature of the crime raises serious issues and concerns in relation to the safeguarding of girls and young women. It is vital that practitioners who come into contact with women, children and families from communities that practise FGM have an adequate knowledge and understanding of the issues in order to respond appropriately and act within contemporary legal frameworks.

Since then The Female Genital Mutilation (FGM) Enhanced Dataset established in 2016 has been recording live data (NHS Digital, 2018 and 2019) to support the Department of Health and Social care's FGM Prevention Programme by presenting a national picture of the prevalence of FGM in England.

It is important to acknowledge that while some health care professionals work closely with communities that have practised FGM for generations, others may rarely come across this practice. Nevertheless, it is important everyone has some understanding of FGM in order to provide the best quality care for the women and girls they come into contact with. This will need to be managed in the context of local safeguarding procedures, which all practitioners should be familiar with.

This publication aims to raise awareness among nurses, midwives, nursing associates and related health care personnel about FGM, and to provide insight and understanding of the socio-cultural, legal and health issues surrounding the practice. Raising awareness is primarily about protecting and supporting girls and women.

Whilst using the terms females, women and girls throughout this document, the RCN also acknowledges that in our gender diverse society, some people who do not necessarily identify as female, may also be at risk of, or survivors of FGM.

Reasons why FGM is practised

A systematic review of global research from El-Dirani et al., (2022) identifies that girls from families whose parents have lower education, are living in rural areas and practising religious beliefs are particularly at risk.

The World Health Organization (WHO) has described FGM as a practice that “reflects a deep-rooted inequality between the sexes, and constitutes an extreme form of discrimination against women”.

FGM is related to the control of women’s sexuality and gender-based social norms relating to ‘marriageability’. FGM is culturally-embedded, as it is viewed as a form of cultural expression among those who support it. FGM may be upheld as a religious obligation by some Muslim populations, even though the practice predates Islam and it is practised by Muslims, Christians and followers of traditional African religions.

In the UK, reasons for practising FGM may have adapted to their context, for instance, the use of FGM to curb sexuality and to preserve girls’ cultural identity, even as prevention of FGM in the country of origin gains ground. Parents may also come under pressure from family and community members in the UK or abroad to have FGM performed on their girls, and need support to avert this (RCM, 2015).

Alongside an overview of FGM and the potential harm and consequences it poses for young women, this guidance provides an outline of the context in which FGM is being managed across the UK. The guidance also provides:

- a consideration of legal and professional requirements, including the mandatory duty to report cases of FGM in those under 18 years old, safeguarding and the importance of multi-agency working
- the national (England only) data set
- clarification of individual nursing and midwifery roles, and a consideration of key service provision requirements
- a review of the practice issues nurses and midwives need to understand.

What is FGM?

The WHO defines FGM as:

“... all procedures that involve partial or total removal of the external female genitalia, or other injury to the female genital organs for non-medical reasons.”

(WHO, 2018)

The practice has no health benefits and can often lead to serious morbidity and mortality in girls and women. The lasting physical and psychological trauma caused by FGM can have lifelong complications for the health and wellbeing of women.

FGM is also referred to as female circumcision, an expression which implies the practice is similar to male circumcision. However, the degree of cutting is far more extensive and the procedure often impairs a woman's sexual and reproductive functions and ability to pass urine normally.

Female genital cutting is another term used to describe the practice, and many traditional societies have different names for what is defined by WHO as FGM (see [Appendix 2: Alternative terms used for FGM](#)).

When discussing FGM, it is important to understand that many women may not be aware of what has happened to them, and because of the use of a variety of local cultural terms (which often refer to purification) may not comprehend what is meant or being referred to when using certain terminology.

Historical and cultural context

The government's strategy for tackling violence against women and girls published in 2021, identified that women and girls affected by FGM may also be at risk from other forms of violence which include so called honour-based abuse – a wide range of abusive behaviours perpetrated by a victim's family or community intended to coerce the victim to behaving to meet family norms or to punish as retribution for a perceived breach of family /community code of behaviours. It includes intimidation, coercion, physical, sexual or psychological violence.

Forced marriage (FM) is where one or both people either do not or cannot consent to a marriage that they are coerced into, and is illegal in the UK.

In 2022, the Health and Care act made the practices of virginity testing (examination of genitalia to determine if intercourse has occurred) and hymenoplasty/hymanoplasty (surgical reconstruction of the hymen) illegal and also highlighted the association between these practices and FGM.

It is important that nurses are aware of the prevalence of these practices and are equipped to safeguard women and girls and offer appropriate support.

Further guidance may be found at: rcnlearn.rcn.org.uk/Search/female-genital-mutilation

Those who do not practise FGM generally view it as a form of abuse and violence and a clear violation of human rights. This attitude is enshrined in numerous international conventions, and agencies, human rights groups, women's groups and governments around the world are committed to eradicating the practice globally.

However, FGM remains deeply rooted within some cultures and traditions, and it can be challenging to rationalise the beliefs that provide a vehicle for the practice to continue. Some communities view FGM as an act of love or a rite of passage and find it difficult to understand why the practice is condemned, believing they are doing the best for their daughters.

It is unclear when and where FGM first started, but reference to the procedure has been found on an Egyptian sarcophagus dating from around two thousand years ago. The practice of FGM was also documented in Britain, Canada and the USA during the 19th century when it was employed to prevent masturbation, cure hysteria and some psychiatric conditions in women (RCN, 2006).

The justifications for performing FGM are many and vary widely between individual communities, contexts and countries; in summary, motivations often relate to the control of women and their sexuality, religion, rites of passage, ideas of hygiene, femininity and aesthetics, as well as social pressures and expectations.

FGM is often erroneously linked to Islam, possibly because it is practised in communities where this religion predominates. Some Muslims believe Islam demands the practice to ensure spiritual purity, but many Islamic scholars disagree and state that there is no reference to FGM in the Qur'an or the Hadith.

In 2014, the Muslim Council of Britain (MCB), the country's largest Muslim organisation, condemned FGM and for the first time issued explicit guidance which criticises the practice and makes it clear it is not supported by any religious doctrine or linked to the teaching of Islam (MCB, 2014).

In reality, FGM transcends religious, racial and social boundaries; a minority of followers of a variety of faiths including Christians, Animists and Jews (the Falashe Jews of Ethiopia) practise it (RCN, 2006).

Performing FGM is seen by some as an essential part of their culture that must be preserved. FGM is often related to ideas about female chastity, hygiene and aesthetics, and is founded on deeply held cultural and traditional belief systems. Illiteracy, the low status of women, their lack of access to money and limited knowledge and power all help to perpetuate FGM.

In some societies FGM is believed to reduce the possibility of premarital and extra-marital sex, improving both the marriageability of 'circumcised' young women and increasing their dowries. Hence, in many places it is viewed as a prerequisite for marriage, which may be the only secure future for women in these societies. FGM may also be considered to promote or maintain virginity and chastity by decreasing women's sexual enjoyment and desire for sex, as well as enhancing their partners' or husbands' pleasure.

The suggestion is that a closed introitus (the opening to the vagina) – also known as infibulation – in which the labia majora and sometimes the labia minora may be sewn together, is considered to provide evidence of virginity. Families therefore view FGM – and by implication, virginity – as important for maintaining their honour in society. The emphasis on ‘tightness’ may be so strong that women wish to be closed again after childbirth, or prior to remarrying if widowed or divorced. FGM may also be falsely believed to improve fertility. It should also be acknowledged that the procedure often carries high social values.

In 2008, a global interagency statement condemned the practice of FGM, including its practice by health professionals:

“Trained health professionals who perform female genital mutilation are violating girls’ and women’s right to life, right to physical integrity, and right to health.”

(WHO, 2012)

Performing FGM is also related to ideas of femininity and masculinity, particularly when the clitoris is likened to a penis. Some communities believe that children are born with the attributes of both sexes and that it is important to ensure that a child is assigned to the appropriate sex and gender role after birth. Therefore, boys must have all feminine attributes removed – the foreskin, which is believed to be the remnants of the labia – while girls must have all masculine features removed – the clitoris, which is believed to be a diminutive penis. These acts are believed to ensure that each child has an unambiguous place in the society.

Aesthetics and cleanliness are other reasons put forward for performing FGM. The female genitalia may be believed to be ritually unclean or polluted; it may also be supposed that a woman’s clitoris ‘poisons’ the baby as it is born. Some cultures see uncircumcised women as bringing shame onto their families; in such societies uncircumcised women, and even girls, may be ostracised, so mothers have strong incentives to make sure their daughters undergo FGM. Many communities view FGM as a positive and normal part of their heritage and perceive continuation of the practice as an aspect of their group identity.

It is also important to recognise that the livelihood of those who carry out FGM for their communities depends on its continuance, so resistance to change may be strong. Such practitioners may also be highly respected members of society. Equally, it must be acknowledged that some health care professionals in the UK come from these communities and may have conflicting views and beliefs about FGM.

Parents who choose to refuse FGM for their daughters may come under considerable pressure from family members to conform. There is a very real fear that despite their objections, elders in the extended family will override their wishes and subject their daughters to FGM.

Prevalence

It is important to acknowledge that available data on FGM is estimated. Global statistics provided by the World Health Organization (WHO, 2023) suggest that FGM is on the decline, however it remains a significant threat to many girls across the globe and is

deeply rooted in regions of Africa, Asia and the Middle East (RCM et al., 2013). The movement of people seeking refuge and asylum from the Horn of Africa region has led to the situation now being taken seriously in the UK. Appendix 1 maps the global prevalence of FGM.

In 2023, the United Nations International Children's Emergency Fund (UNICEF) estimated that at least 200 million girls and women in 31 countries have been subjected to FGM across Africa and the Middle East. These are regions where FGM is primary concentrated, and these women are living with the consequences of having had FGM performed on them. For example, UNICEF (2019) reports that:

‘FGM is nearly universal among girls and women of reproductive age in Egypt.’

It should also be understood that some Asian countries also practice FGM, such as Indonesia, where approximately half of girls under the age of 12 have undergone some form of FGM (UNICEF, 2019).

While not all women are negatively affected by FGM, research demonstrates that large numbers are physically and psychologically damaged by this mutilation (WHO, 2023). These numbers do not account for those who have died as a direct or indirect result of having FGM performed on them, or the babies that have died following a traumatic childbirth due to complications from FGM.

WHO (2023) also reports on worrying evidence that the average age at which the practice is carried out is falling in some countries. One of the cited reasons for this shift includes the wish to have girls cut more discretely (or underground), in the presence of legislation against FGM. Another possible adverse effect of legislation is its tendency to encourage a cross-border movement of women from a country where the practice is illegal to another country where it is allowed.

Many girls and women die from the short-term effects of FGM which include haemorrhage, shock or infection, whilst significantly more suffer lifelong disability and may die from longterm effects such as recurrent urinary or vaginal infections. Pain during intercourse and infertility are further common consequences of FGM.

FGM increases the risk of women dying during childbirth and makes it more likely that the baby will be born dead. This increased risk can be as a result of severe bleeding and obstructed labour in places where safe and appropriate maternal health services are inadequate or inaccessible. In Somalia, where 90-98 percent of women are infibulated, one in every 100 women giving birth dies as a result of this procedure (see Figure 1 for a short narrative that expands on this point).

The WHO, which heads up the global initiative against FGM, published a strategy to counter the medicalisation of FGM entitled *Global Strategy to Stop Health Care Providers from Performing Female Genital Mutilation* (WHO, 2014), which has formed the basis of work by WHO since then. The strategy was developed in collaboration with other key UN agencies and international organisations following the adoption of a resolution to eliminate FGM by the UN General Assembly, which was passed in December 2012.

This resolution resulted in greater engagement by global organisations such as WHO, UNICEF and the UNFPA, as well as the European Parliament and other national governments, in working to end this violation of the human rights of girls and women.

In 2009, END FGM, a European campaign led by Amnesty International Ireland, began working in partnership with a number of organisations in the European Union (EU) to highlight increasing concerns about the prevalence of FGM across the EU (endfgm.eu). The campaign focused on human rights issues and the need to lobby for a comprehensive and coherent approach towards ending FGM (END FGM, 2009). In 2023, the End FGM campaign, who produce an interactive map on their website (endfgm.eu) estimated that there are approximately 600.000 women living with the consequences of FGM in Europe.

Tackling FGM in the UK

The last decade has seen a noticeable increase in activity focused on the elimination of FGM because of the devastating consequences to women has been made possible, in part, by the bravery of some girls and women who have been prepared to share their experiences and risk their own wellbeing in order to campaign against this practice.

A study of one such community-based project by Brown and Hemmings (2013), demonstrated how programmes which focus on community-orientated strategies can achieve significant outcomes by engaging with community members and leaders to change attitudes and deliver education on the risks associated with FGM. The evidence shows that understanding the community and providing access to specialist FGM services are vital steps for changing the existing culture.

Key health care professionals such as school nurses and health visitors have identified roles in recognising vulnerable families, and establishing relationship with those who may be at risk, which should also help to reduce the risk of harm, if managed in a sensitive but supportive environment.

Hussein's study (Hussein, 2010) also provided useful insights into women's experiences, perceptions and attitudes to FGM, and highlights key actions that can help eliminate FGM:

- engagement with the community and integration
- providing a safe space for dialogue and discussion
- raising awareness of FGM
- providing specialist health services which should incorporate female practitioners, interpreters, physical and psychological support, and sensitivity by staff in understanding FGM
- awareness of the law and balanced supportive safeguarding frameworks.

There are numerous examples of programmes across the UK, some initiated by women and others by professionals working across agencies who have invested in campaigns to raise consciousness and understanding of the consequences of FGM. These include community groups and the police, as well as education and health and social care practitioners. forwarduk.org.uk.

The education of male partners and community leaders may also reduce the number of children, and young and older women who suffer in the future. However, cultural practices such as FGM have been ingrained for many generations and require extensive education to address the issues thoroughly and effectively.

The establishment of an FGM Hotline (0800 028 3550 or fgmhelp@nspcc.org.uk) which is managed by the National Society for the Prevention of Cruelty to Children (NSPCC), has proved a useful source of support for women and professionals.

In 2013, the Royal College of Nursing, the Royal College of Midwives, the Royal College of Obstetrics and Gynaecology, the Community Practitioners and Health Visitors Association, in association with the human rights organisation Equality Now, created an intercollegiate FGM group (RCM et al., 2013) to develop recommendations towards tackling FGM across the UK, many of which are now being implemented.

The intercollegiate group's recommendations clearly assert the view that FGM is child abuse and a violation of human rights and provides an outline of areas for service improvement (see Figure 2). The report called on health care professionals, together with the police and education and social work professionals, to consider their responsibilities on the safeguarding of the girls and women who may be affected by, or at risk of being mutilated.

Figure 2 – Intercollegiate recommendations for tackling FGM in the UK

1. Treat it as child abuse.
2. Document and collect information.
3. Share that information systematically.
4. Empower frontline professionals.
5. Identify girls at risk and refer them as part of child safeguarding obligations.
6. Report cases of FGM.
7. Hold frontline professionals accountable.
8. Empower and support affected girls and young women (both those at risk and survivors).
9. Implement awareness campaign.

(RCM et al., 2013)

There are a number of immediate challenges when it comes to tackling FGM; these are related to the education of the public and professional workers, the absolute need to work collaboratively with communities, and the need for more accurate data on actual prevalence. High quality information-sharing pathways across agencies and comprehensive evidence gathering to support prosecutions, where a child has been put at risk or mutilated (RCM et al., 2013) is also essential.

Since the publication of these recommendations a wide range of initiatives have been enhanced or established with the intention of tackling FGM across the UK and the globe. Many of these recommendations have in total or in part been implemented and remain key issues for tackling FGM across the UK.

The most notable changes from a health care practice perspective have been the collection of anonymised data via the Enhanced Data Set and the mandatory duty to report all cases of FGM in those under 18 years of age.

Effective use of data

It is critically important to have accurate data on the actual numbers of women affected, by which types of FGM. In 2014, as part of its FGM Prevention Programme, the Department of Health and Social care (ISB, 2014) set in place a requirement for better data collection across the health service and the first anonymised data report was published by the HSCIC (Health and Social Care Information Centre, known as NHS Digital) in October 2014. In February 2023, NHS England and NHS Digital merged.

Enhanced Data Set is the term used for the collection of data on all cases of FGM, which is published anonymously by HSCIC. It is currently collected from acute trusts, mental health services and GP practices. The main purpose is to collect more accurate data on the numbers of women affected by FGM, and their location, in order to be able to ensure appropriate services are commissioned. This has not been implemented without concerns about confidentiality, and whether it might put women off seeking the health-related services they need. At present there is no evidence to support this concern, however the emerging picture will continue to be scrutinised locally and nationally. For further information see: gov.uk/government/publications/fgmenhanced-datasetguidance-on-nhs-staffresponsibilities

This only applies to health care where the woman has to use an NHS Number, for example sexual health clinics are currently exempt from this requirement as they do not use NHS registration numbers.

Initial figures show that between April and September 2014, 1,279 female patients previously identified as having been subjected to FGM were accessing health care. This data represents the first official figures published on FGM cases seen in hospitals in England, and should be viewed critically as it will not provide a complete picture for some years to come. This was followed by further reports, available at <https://digital.nhs.uk>, which begins to provide a country-wide profile of women seeking health care who have been subjected to FGM.

This data should support better planning and commissioning of services; especially in relation to identifying where services are most needed.

The mandatory duty to report FGM cases to the police came into effect from 31 October 2015. The new duty applies where a nurse, midwife or nursing associate (and extends beyond health care), in the course of their work, is either informed directly by the girl that FGM has been carried out on her, or observes physical signs which appear to show that she has been abused by FGM having been carried out and has no reason to believe that the act was necessary for the girl's physical or mental health or for purposes connected with labour or birth.

The duty applies only to girls who are under 18 at the time of disclosure or visual identification of FGM by health care professionals and applies only in England and Wales.

The mandatory duty to report FGM requires health care professionals to report all cases of FGM in under 18-year-olds directly to the police, as well as implementing a

safeguarding pathway for the girl, and ensuring she receives the required care. The Department of Health and Social care has provided details of how the process should be implemented locally, however it is critically important that all nursing and midwifery staff familiarise themselves with local procedures, and contacts for safeguarding pathways.

Appendix 3 provides an overview of the process outlined by the Department of Health and Social care and NHS England, endorsed by a number of Royal Colleges.

Scottish Government: Violence against women and girls (VAWG) gov.scot/policies/violence-against-women-and-girls/female-genital-mutilation-fgm/

National Safeguarding Team (NHS Wales): wales.nhs.uk/sitesplus/documents/888/FGM%20Annual%20report%202017-18%20Final.pdf

NHS England. Commissioning services to meet the needs of women and girls with Female Genital Mutilation: england.nhs.uk/publication/commissioning-services-to-meet-the-needs-of-women-and-girls-with-female-genital-mutilation-fgm

In Northern Ireland further information can be found at Safeguarding Board for NI – FGM: safeguardingni.org/children-and-young-people/advice-parents-and-carers/female-genital-mutilation-advice

In complying with the NMC code, nurses and midwives should continue to have regard to their wider safeguarding responsibilities, whether in relation to FGM or any other forms of abuse.

Nurses, midwives and nursing associates should familiarise themselves with the local safeguarding processes which should now include the pathway for best supporting a girl under 18 years of age found to have been abused in this way. A range of additional resources to support health professionals comply with the new duty are available at: gov.uk/government/publications/mandatory-reporting-of-female-genital-mutilation-procedural-information. These resources include a DHSE risk assessment tool, and specific safeguarding guidance for professionals. Failure to comply with this duty to report may result in an investigation of the nurse or midwife's fitness to practise by the NMC.

FGM is a safeguarding issue and the main difference between discovering or suspecting FGM has been performed on a girl and any other form of abuse is that the practitioner has to report it directly to the police. This is being managed locally through the non-emergency 101 service (police.uk/pu/contact-us), and any nurse or midwife who is likely to provide care for girls, should be familiar with these local systems. The DHSE has provided guidance on what information should be provided and this is available at: gov.uk.

At the same time they should implement their local safeguarding processes as they would for any form of suspected or confirmed abuse, as well as providing care for the initial health query.

At present there is no requirement to refer women (ie, over 18 years of age) with FGM to social services or the police unless they have specific safeguarding concerns, however there should be opportunities to refer individuals to specialist health services where complications are identified or where other safeguarding concerns have been identified for example other child at risk, or other vulnerabilities. Health professionals need to

be aware of which services are available locally to support women who have physical, psychological or psychosexual complications from having had FGM performed on them.

Government initiatives include the development of multi-agency guidelines to support best practice (HM Government, 2016) and in February 2014 the UK government published its declaration to end FGM on the International Day of Zero Tolerance to Female Genital Mutilation (6 February 2014). A national FGM prevention programme, initiated by the Department of Health and Social care in England, is well underway and [Appendix 4](#) provides a diagrammatic overview of the government departments currently engaged in the campaign to end FGM across England.

The UK government has amended the existing Female Genital Mutilation Act 2003 so that prosecutions can be made to prevent British citizens and residents living in this country from taking children overseas for the procedure. There are a series of amendments included in the Serious Crime Act 2015, which strengthen the legal framework regarding FGM. It means that the 2003 Act can capture offences of FGM committed abroad against UK residents, irrespective of immigration status. The new law also allows for civil protection orders to be made where a girl or woman is identified at risk of FGM.

Scotland

In Scotland, the FGM programme is managed under the Violence Against Women Outcomes Framework (further information, including fact sheets in a range of languages, can be found at gov.scot). A critical range of stakeholders are engaged in developing services, including data collection, learning about FGM and community engagement, as well as an initiative to engage head teachers, where educational establishments are seen as key players in protecting girls who may be at risk.

In January 2016, Scotland consulted on new guidelines and in 2017 published a framework for agencies and practitioners to develop and agree processes that promote the safety and wellbeing of women and girls. This is available at: gov.scot.

Wales

The Welsh government has an ongoing programme of work developing a national training framework for key professionals across the public sector; including writing to all schools on the subject. In 2019, the Welsh government published its guidance for professionals, available at: gov.wales.

Northern Ireland

In Northern Ireland, the FGM programme of work is managed within safeguarding and the Department of Health, Social Services and Public Safety for Northern Ireland (DHSSPSNI) has published guidelines (DHSSPSNI, 2014). These guidelines provide a resource for contacts and specific guidance for Northern Ireland.

There is growing expertise and understanding of the need to enhance services for women with FGM and key organisations are working to enhance service provision and awareness raising.

The FGM mandatory reporting duty applies in England and Wales only. For health care professionals working in Scotland or Northern Ireland, existing safeguarding responsibilities should be adhered to.

The e-FGM educational programme

This e learning module was developed for healthcare and is provided free to all healthcare professionals, including school nurses, practice nurses, health visitors and GPs. e-lfh.org.uk/programmes/female-genital-mutilation.

NHS England national FGM support clinics for non-pregnant women

In September 2019, several community satellite clinics across England were launched specifically to provide specialist support for non-pregnant survivors of FGM. These clinics are the first of their kind and offer a wide range of holistic care. Operating on both a walk-in and referral basis, women over 18 can attend to access care for physical, mental, emotional and social wellbeing. Clinics are led by an all-female team of specialist midwives and nurses, supported by FGM advocates and counsellors, with clearly defined pathways to referral to tertiary care where required. Women may access medical advice and treatment in relation to the effects of FGM, including assessment, routine sexually transmitted infection screening, cervical screening, access to deinfibulation and referral to urology and/or gynaecology services where appropriate. The staff at the clinics will consider the safeguarding needs of clients and will work closely with local multi-agency partners. They have been set up with health care trusts which have been identified through the NHS enhanced data set as reporting the highest incidence of patients with FGM and which already have significant experience in providing specialist services. The effectiveness of this initiative will be subject to close evaluation in terms of its ability to impact physical and mental health, social and emotional wellbeing and its contribution to working with local communities to support survivors and their families.

More information on the clinics is available at: nhs.uk/conditions/female-genital-mutilation-fgm/national-fgm-support-clinics

Global commitment to end FGM

In 2016, the United Nations launched 17 goals for global sustainable development by 2030, aimed at creating a future that eradicates poverty, promotes social inclusion and protects and sustains the environment. Goal 5 promotes gender equality which cannot be achieved without eliminating violence against women and girls in all its forms. It is estimated that 68 million women and girls worldwide are at risk of FGM between 2015-2030, and ending this violence is critical in achieving other sustainable goals such as promoting health and wellbeing, achieving safe motherhood, facilitating social inclusivity and enabling equal access to education. Zero Tolerance Day on 6 February each year promotes the campaign to end FGM by focusing on the achievements of work towards this.

More information is available at: un.org/en/events/femalegenitalmutilationday

The Girl Generation is now a global campaign that supports the Africa-led movement to end FGM. Further information is available at: thegirlgeneration.org

FGM procedures and health effects

Female genital mutilation (FGM) is a procedure that involves partial or total removal of the external female genitalia, or causes other injury to the female genital organs for non-medical reasons (WHO, 2018).

FGM may be conducted from shortly after birth to age 15 or young motherhood. The procedure may be performed in a clean, clinical setting but is often undertaken in poor light, without anaesthesia and using blades, knives, broken glass or non-surgical instruments that are often shared.

Girls have to be forcibly restrained and, following more extensive forms of FGM, their legs may be tied together for days to aid healing. Accidental damage, infection and haemorrhage are common, and long-term physical and mental health problems may follow if the child survives – death is not uncommon.

FGM offers no therapeutic benefit to women and girls, and is illegal in many countries including the UK. Some countries have legalised FGM on the assumption that it is safer if conducted within medical care, however this is contrary to WHO recommendations (WHO, 2008).

Types of FGM

FGM is practised in different ways by different communities, and some forms are more extensive than others and cause greater health risks for girls and women. The WHO has categorised FGM into four types (see Figure 3).

Figure 3 – WHO classifications: types 1-4

The WHO has developed four major FGM categories.

1. Clitoridectomy: the partial or total removal of the clitoris – a small, sensitive and erectile part of the female genitals – or the removal of the prepuce only – the fold of skin surrounding the clitoris, also known as the clitoral hood, rarely, if ever performed alone. – See [Diagram 2](#).
2. Excision: the partial or total removal of the clitoris and the labia minora, with or without excision of the labia majora – the labia are the ‘lips’ that surround the vagina) – see [Diagram 3](#).
3. Infibulation: the narrowing of the vaginal opening through the creation of a covering seal. The seal is formed by cutting and repositioning the inner, or outer, labia, with or without removal of the clitoris – see [Diagrams 4 and 5](#).
4. Other: all other harmful procedures to the female genitalia for non-medical purposes; for example pricking, piercing, incising, scraping and cauterising the genital area – see [Diagram 6](#).

Diagram 1 – Normal external female genitalia




Diagram 2 – Type 1 FGM

When the clitoris or the clitoral hood is cut off




Diagram 3 – Type 2 FGM

When the clitoris and inner lips are cut off






Diagram 4 and 5 – Type 3 FGM

When the clitoris, inner lips and outer lips are cut and sewn together

**Diagram 6 – An example of type 4 FGM stretched labia**

All other harmful practices including stretching, pricking, piercing, cutting, scraping and burning



Health risks and complications

The complications that may occur following FGM will depend on the type and extent of the procedure carried out and should not be underestimated. These are generally classified as:

- immediate (Figure 4)
- intermediate (Figure 5)
- long-term complications (Figure 6).

Figure 4 – Immediate complications

Immediate complications include:

- haemorrhage, pain, shock
- wound infection, septicaemia, tetanus
- urine retention
- injury to other tissues, for example, vaginal fistulae
- ulceration of the genital region
- bacterial or viral infections such as Hepatitis and HIV due to instruments being re-used without sterilisation
- death.

Figure 5 – Intermediate complications

Intermediate complications include:

- delayed healing
- abscesses
- scarring/keloid formation, dysmenorrhoea and haematocolpus – obstruction to menstrual flow
- pelvic infections
- obstruction to urinary flow
- urinary tract infection.

Figure 6 – Long-term complications

Long-term complications include:

- psychosocial trauma and flashbacks, post-traumatic stress disorder
- lack of trust in carers
- vaginal closure due to scarring
- epidermal cyst formation
- neuromata – benign tumours of nerve tissue that may arise from cut nerve endings and cause pain
- pain and chronic infection from obstruction to menstrual flow
- recurrent urinary tract infection and renal damage
- painful intercourse (dyspareunia), lack of pleasurable sensations and orgasm, marital conflict
- infertility from pelvic inflammatory disease and obstructed genital tract
- risk of HIV through traumatic intercourse
- childbirth trauma – perineal tears and vaginal fistulae
- postnatal wound infection
- prolonged or obstructed labour from tough scarred perineum, uterine inertia or rupture, and death of infant and mother
- vaginal fistulae as consequence of obstructed labour.

Serious illness and death can occur even when FGM is carried out by health professionals, who may be acting illegally and in unclean surroundings without sterilisation facilities for instruments. Even where the practitioner is skilled and cleanliness ensured, the long-term effects can ruin women's lives and relationships.

Type 3 FGM inevitably causes more health problems and deaths. Most women with type 3 FGM tend to have problems with penetration; for some couples it can take several months to achieve this. Men who find penetration difficult at first intercourse may cut the introitus to make this easier, with pain, infection and bleeding as a consequence for the woman. In some cultures, the introitus will be opened surgically immediately before the first intercourse. The pinhole introitus left after FGM and lack of sexual response following type 3 FGM can cause significant problems for some women and their relationships. The physical difficulties, ongoing dyspareunia, infection, scarring and psychological problems may lead to infertility and consequent rejection of women by some men. The problems can continue for years.

Type 3 FGM can cause particular dangers in childbirth. In addition to the problems listed above, prolonged or obstructed labour and perineal laceration occur due to tough, unyielding scar tissue. Clitoridectomy (type 1 FGM) does not usually cause obstruction unless there was infection at the time of mutilation.

Neonatal problems occur primarily as a result of obstructed or prolonged labour which, if unchecked, can cause fetal distress, anoxia (lack of oxygen to the body's tissues) and fetal death.

The psychosocial trauma and post-traumatic stress caused by FGM cannot be underestimated and it may include fear of sexual intercourse, post-traumatic stress disorder, anxiety, depression and memory loss; its impact is felt on the lives of women and their families. It is an area of practice that is increasingly becoming better understood and researched; a pilot study (Liao LM et al., 2013) demonstrated the critical need for further research in this area, in particular the requirement for psychological and psychosexual research to enhance service provision. It is also becoming more widely acknowledged that FGM has a detrimental impact on a woman's sexual pleasure, as well as her physical/sexual and mental wellbeing.

The emergence of specialist clinics, such as those for children and those catering for the psychosexual needs of women affected by FGM, is a further demonstration of the advances being made in providing better services for women and girls. The fear of sexual intercourse, symptoms of post-traumatic stress disorder, including anxiety and depression should be considered when caring for women who have been subjected to abuse, consequently, service provision should encompass a wide range of services required by women who have been traumatised by FGM – in particular, mental health, psycho-sexual services and counselling. These should be specialised services, and need to be sourced locally to best support women who have been affected.

Human rights, legal and professional responsibility

A human rights issue

FGM is a violation of respect for, and the dignity of, girls and women subjected to this cruel practice; it is a clear form of violence against women and girls. The practice breaches fundamental human rights guaranteed by a multiplicity of international agreements, the most significant of which in terms of UK law is the European Convention for the Protection of Human Rights and Fundamental Freedoms. Drawn up by the Council of Europe in 1950, the Convention has been incorporated into domestic UK law through the Human Rights Act (2000). The Act affords citizens a variety of legal remedies in circumstances where their rights have been interfered with.

Relevant rights in the context of FGM include:

- Article 3 – protection against inhuman or degrading treatment
- Article 8 – the right to respect for privacy and family life.

The requirements of the European Convention reflect, very closely, existing good professional practice. A failure of the state to fulfil its positive obligation to protect child and adult female rights in these circumstances, by prosecution or otherwise, could itself be open to challenge under human rights legislation.

Various forms of mutilation, whether carried out for religious or social reasons, and conducted without the child's consent and for nontherapeutic purposes, infringe the child's right to bodily integrity.

Although parents have rights to bring up their children according to their own beliefs, the rights of the child to protection come first and courts will inevitably weigh the balance more heavily in favour of child protection. There is also a large body of international human rights law specifically to protect and promote the rights of children; for example, the *UN Convention on the Rights of the Child and the Protocol to the African Charter* (African Union, 2003).

Legal aspects

FGM is illegal in a number of countries, even those where it is customarily practised. Many countries where FGM is not normally carried out, such as the UK, also have legal provision to cover those who arrive from elsewhere, especially if they are migrants from FGM practising communities. This includes protection for those temporarily removed from the UK with the intention of inflicting harm on them.

The UK legal framework

The UK Acts of Parliament are relevant in respect to FGM:

- Prohibition of Female Circumcision Act (1985)
- Female Genital Mutilation Act (2003)
- Serious Crime Act 2015.

The 2003 Act applies to England, Wales and Northern Ireland; in Scotland, the Prohibition of Female Genital Mutilation (Scotland) Act 2005 applies. Doctors, nurses and midwives participating in FGM also face removal from their respective professional registers and would be prosecuted for taking part.

The 1985 Act states that it is an offence for any person to:

- ‘... excise, infibulate or otherwise mutilate the whole or any part of the labia majora or clitoris of another person’
- ‘... aid, abet, counsel or procure the performance by another person of any of those acts on that other person’s own body’; this also means that following childbirth or de-infibulation, the anterior middle incision can only be over-sewn and not closed back to its original state.

However, because of concerns relating to girls being taken out of the country for FGM and the lack of prosecutions, it became necessary to amend the law and repeal the 1985 Act. The Female Genital Mutilation Act (2003) came into force in March 2004. It re-enacted the above offences and created additional new offences, sending a strong message to communities practising FGM and practitioners involved in aiding, abetting, and/or counselling to procure, and performing FGM, that the practice is no longer acceptable in the UK even if performed in another country.

The main changes made were:

- it was now against the law for UK nationals or permanent UK residents to carry out FGM abroad, or to aid, abet, counsel or procure the carrying out of FGM abroad, on a UK national or permanent UK resident, even in countries where the practice is now legal (Female Genital Mutilation Act, 2003). This means that the law protects any girl, who is a UK national or permanent resident, from FGM anywhere in the world
- the penalty was increased from five to 14 years’ imprisonment
- the term ‘mutilation’ is used instead of circumcision
- the term ‘girl’ includes ‘woman’
- aiding, abetting and counselling applies to those who assist or persuade a girl to perform FGM on herself even though it is not itself an offence for that child to carry it out on herself (although not an offence, consideration should be given to whether such self-harm is a safeguarding issue, especially where the action may be the result of adult pressure)
- it is now considered illegal to re-infibulate a woman following the birth of her baby, because this is seen as a form of FGM, within the WHO definition of FGM.

Despite this robust legislation, the intercollegiate report (RCM et al., 2013) considered the lack of prosecutions to date; there is evidence that the link between prosecutions and prevention is becoming increasingly recognised. There are renewed efforts among the police and others, including health professionals, to directly address the barriers to prosecution.

Following extensive evidence gathering, in July 2014 the Home Affairs Committee published its report on FGM (House of Commons Home Affairs Committee, 2014) which called for a national action plan and highlighted a number of issues relating to:

- impediments to achieving prosecutions for offences against the law on FGM
- the need to revise the multi-agency guidelines on FGM to include the mandatory questioning on FGM for antenatal booking interviews and at GP registration, and a requirement for the personal child health record (also known as the PCHR or 'red book') to refer explicitly to FGM (House of Commons Home Affairs Committee, 2014)
- requiring all schools to provide training on FGM
- changes to the law on FGM, including the creation of FGM protection orders (similar to those for forced marriage).
- that the failure to report child abuse should become a criminal offence if other measures to increase the level of reporting are not effective in the next 12 months.
- specialist services for women and girls living with FGM.

This was contemporaneously published with a Ministry of Justice consultation on whether it was necessary to have a specific civil protection order, akin to the forced marriage protection order, that might provide an additional tool to prevent female genital mutilation and which could complement the existing criminal law (MoJ, 2014); this could also provide some anonymity for those giving evidence. In October 2014, the Ministry of Justice announced it would be implementing protection orders which came into being via the Serious Crime Act in 2015.

The Serious Crime Act (2015) made provision to extend the jurisdiction of the Female Genital Mutilation Act (2003). The changes mean the 2003 Act will capture FGM offences committed abroad by or against those habitually resident in the UK, irrespective of whether they are subject to immigration restrictions. Provided the offence is committed at a time when the accused person and/or the victim is resident in this country, it should not matter whether or not they intend to live here indefinitely or whether they also live elsewhere. See also the mandatory duty to report (on [page 13](#)).

Safeguarding now and for future generations

All health care professionals have a duty of care to girls and women at risk of having FGM performed, or who have been cut in the past. Their responsibilities include ensuring their practice is performed within the requirements of their regulators (for example, the NMC for nurses, registered nursing associates and midwives) and the overall legal framework of the country they practice in.

While the overarching legal issue related to FGM is its illegality, practitioners must also ensure they provide care and support that is consistent with safeguarding law and procedures. Professionals should be familiar with what to do if they are worried that a child at risk of and/or being abused, including the local child protection policies.

The current child protection policies in UK countries are as follows:

- England – *Working Together to Safeguard Children* (2018) workingtogetheronline.co.uk
- Scotland – *National Guidance for Child Protection in Scotland* (2014)
- Northern Ireland – *Co-operating to Safeguard Children and Young People in Northern Ireland* (2017)
- Wales – *All Wales Child Protection Procedures Review Group* (2018).

Safeguarding law provides the framework through which a girl or woman's needs are assessed and her best interests considered. The welfare of the child is paramount, according to the Children Act (1989) which allows legal action to be taken. However, legal measures may not be appropriate if protection can be achieved without these; judgemental attitudes are potentially harmful and bringing about change is more effective if people's long-held attitudes are addressed.

It is important to promote understanding and to protect girls and women from the practice through a continuing programme of education and raising awareness. This needs to include explaining why FGM is considered to be a violation of human rights, and the connection between the procedures and the long-term effects on the body, psychological wellbeing and emotions.

All regions and countries in the UK have in place procedures for safeguarding children, young people and vulnerable adults, and all practitioners must ensure they are confident in using them in their practice.

The NMC website (nmc-uk.org) contains safeguarding guidance and information on how to raise and escalate concerns and apply safeguarding principles in nursing and midwifery practice. Similarly, the UK government's Children's Services provides online information and resources on safeguarding (gov.uk) including Working Together to Safeguard Children, available at: gov.uk/government/publications/working-togetherto-safeguard-children--2, and the RCN has produced guidelines on Safeguarding Children and Young People: Roles and Competencies for Healthcare Staff (2019) which is available at: rcn.org.uk/publications.

Safeguarding girls at risk of harm through FGM poses specific challenges because the families may give no other cause for concern (such as parenting responsibilities or relationships with their children). Family members may believe FGM is the right thing to do and consider it is in the child's best interest, and adults may find it difficult to understand why the authorities should intervene in what they see as a cultural practice specific to their way of life. The family situation may be compounded by those who wish girls to be ritually cut when others disagree. Similarly there may be an inter-generational element, or a mother and father may have differing views about their daughters.

The desire to carry out FGM is also not confined to individuals within particular levels of education or social class. The pressure to undertake this procedure may be embedded in family structures. At all times it is important to 'think the unthinkable', and act with 'respectful uncertainty' (DH, 2003).

Four specific issues are important in this context:

1. FGM is an illegal act; regardless of their age, a girl or woman has the right to protection from activities or events that may cause her harm. These rights (UN Convention on the Rights of the Child) are enshrined in UK human rights legislation and is reflected in other laws, including the Children Act 2004 (England and Wales), Protection of Children Act (Scotland) 2003 and the Children (Northern Ireland) Order 1995. These rights and protections are in addition to other legislation that criminalises the practice of FGM.
2. The need to safeguard children and young people involved in FGM.
3. The risk to girls where a related adult has undergone FGM.

4. Situations where a child may be removed from the country for the purposes of performing FGM. Taking a girl abroad to perform FGM is illegal; however, there may be instances where the exact risk is not known but one parent may be concerned enough to alert professionals. In certain circumstances the Child Abduction and Custody Act (1985) can be used to prevent a girl being removed from the country. This legislation has a requirement for both (married) parents to agree to a child leaving the country. Normally a prohibitive steps order made by social services will suffice.

Whenever there is concern that a girl is at risk of harm through FGM, steps must be taken to safeguard them. This has to take account of the mandatory duty to report all cases to the police, as well as implementing local safeguarding procedures. If she has already had the procedure performed and there are other female siblings in the family, a child in need referral should be made following the steps outlined in *What to do if you're Worried a Child is Being Abused* (HM Government, 2015).

The referring practitioner should follow guidelines about working in partnership with the family by being honest where this is possible and handling any disclosure sensitively. But the practitioner must also be clear about the reasons why they are undertaking safeguarding actions. This partnership would be unacceptable where the girl may come to harm as a result of any evidence being given to parents, as it could cause the family to vanish with their daughter.

There may be a need to approach social services first with suspicions. As well as health professionals having a role in providing information, it is the responsibility of social services to provide the family or parents with information about UK law and policy around FGM, safeguarding and support mechanisms. Social services should also alert families to their right to seek independent legal advice should they wish to appeal against any specific interventions.

The Department of Health and Social care has a new framework – developed in conjunction with professional bodies, Royal Colleges and the Department for Education – published to guide professionals dealing with girls at risk of FGM from birth onwards, so that staff will know how to respond to FGM safeguarding concerns. *Female Genital Mutilation Risk and Safeguarding Guidance for Professionals* (DHSE, 2017) includes a risk assessment process and can be found at gov.uk. The information is being regularly updated.

All professionals coming into contact with children and young people have a responsibility to safeguard and promote their welfare and should know what to do if they have concerns about safeguarding issues, including child protection. The publication *Safeguarding Children and Young People: Roles and Competences for Health Care Staff* (RCPCH) provides extensive details on competences, skills and attitudes which should be embedded in all levels of practice and contact around safeguarding.

The DHSE has also produced a range of resources, including a DHSE Risk Assessment tool, and specific safeguarding guidance for professionals (*Female Genital Mutilation Risk and Safeguarding Guidance for Professionals*) available at: gov.uk

Across the UK, specialist safeguarding/child protection professionals provide expertise and have specific roles and responsibilities in safeguarding children. In England, Northern Ireland and Wales, named and designated professionals perform this function while in Scotland nurse consultants, child protection advisers and lead clinicians fulfil specialist roles.

Service provision and multi-agency working

Communication

Communication with women, even if interpreters are not required, needs to be clear, using straightforward language and explanations. Pictures or diagrams may help. It is important to listen without interruption, avoid rushing or providing too much information at once, and check that women have understood. Some of these women may not have seen female genitalia which has not been mutilated therefore it may be useful to show diagrams of both for comparison.

Initiating the conversation can be challenging for many health care practitioners and so framing it in the context of “have you ever been cut or had any form of surgery or piercings?” may be a useful opening question to encourage further discussion.

There are a range of films available about having conversations about FGM which is available at: gov.uk/government/publications/female-genital-mutilation-resource-pack/female-genital-mutilation-resource-pack#dvds-and-videos

A Statement Opposing Female Genital Mutilation, published in June 2015, can be given to girls and families to help them when explaining the illegality and complications of FGM to wider family and community members, both when they are abroad and in the UK, and is available at HM Government – Health Passport (Statement opposing female genital mutilation).

Multi-agency working

While raising awareness and creating cognisance of FGM are critical in the fight to eradicate the procedure, there also needs to be adequate and well considered services to support best practice; whether that is safeguarding, data collection, physical and psychological care, and training and education. A key characteristic of high quality safeguarding is multi-agency working and managing FGM similarly depends on this; work needs to reflect local need, and extend across health and social care, as well as education and the police forces.

This publication is focused on the needs of nurses and midwives, but providing the best care for women and girls affected by FGM must include working alongside other agencies involved in safeguarding. Safeguarding is everybody’s business and should not be left to those who may lead services in this crucial area of work.

FGM is an issue where multi-agency teamwork and communication is vital. All services should be open, offer flexible access and there should be cohesive collaboration between agencies. Women may be unwilling to come forward for help, or may be unaware of what is available, or not know how to ask; they may find it difficult to raise the topic with health care staff because they may believe practitioners have limited awareness of FGM or may respond in a negative manner. Nurses and midwives should be alert to this and take opportunities to enquire sensitively and offer support and referral to specialist clinics. Generally, women are likely to prefer female carers to male.

It is important to note that health care professionals may not need to provide all services.

Support groups and organisations have a very important role to play and have been prime movers in bringing about change.

Service provision

A seamless service for women and adequate protection for girls at risk depends on integrated working between services. Everyone who may come into contact with FGM practising communities needs to understand their responsibilities, and have appropriate training and referral mechanisms; they will need to know from whom they should seek help and advice and the steps to take to provide appropriate support. This means clear, well understood and rehearsed guidelines must be in place. Similarly, other care practitioners must be aware of FGM issues and be able to recognise when girls or young women may be at risk or have been harmed.

Health care managers and commissioners should focus on access to and provision of:

- clear guidance to employees
- education and training for all, preferably across agencies, as well as specific to health care
- clear lines of communication with others such as education, social and law enforcement services.

The DHSE document *Commissioning Services to Support Women and Girls with FGM* (Department of Health and Social care, 2015) sets out what a successful and safe service to support women and girls with FGM might look like.

Many of the requirements for women's primary, sexual and reproductive health services, such as routine urinalysis, cervical screening, gynaecological and fertility services, may not be well-prepared to meet the needs of women and girls who have been subjected to FGM, and may need further support and changes to established service delivery.

Women and children who have been mutilated may need access to services such as:

- counselling, psycho-sexual and psychiatric support through statutory or voluntary services because of psychological trauma, relationship or psycho-sexual difficulties and/or support with contraception
- assisted conception/fertility clinics
- uro-gynaecological services, including surgical reversal of infibulation
- access to an interpreter service with workers who appreciate the problems facing children and women who have been cut, and also those of refugees and asylum seekers: it is essential that women are not reliant on family members for interpretation when dealing with health care professionals; children should never be used for interpreting purposes
- specialised maternity advice services.

It is also important for women and girls to have access to specialist services. A growing number of specialist clinics are becoming available across the UK and all nurses and midwives, particularly those working with women and children, families and communities, should have the appropriate specialist skills to work effectively with this client group.

Health visitors, school and community children's nurses

Health visitors, school and community children's nurses (CCNs) have a responsibility to ensure families know that the practice of FGM is illegal and are in an ideal position to act if they consider a girl or young woman is at risk. FGM may be carried out secretly in the UK, but it is more likely that a girl or young woman will be sent 'home' to her family's country of origin for FGM to be performed. This is inevitably, although not exclusively, likely to occur during a school holiday, although approaches will vary from one community or ethnic population to another.

It is therefore essential that those coming into contact with girls and young women have detailed knowledge of local communities and social structures, whilst engaging with them to really understand what is important to them, as well as being aware of the safeguarding responsibility.

Health visitors work closely with families in their homes and have a key role to play in health promotion and education from an early age in a girl's life. This may include helping and supporting families to explore ways of breaking the cycle of ritual abuse. Health visitors, school nurses and CCNs are also well placed to collaborate and engage in support and referral as part of a multi-professional team. School nurses and CCNs, like teachers, may be in a position of trust and receive disclosures from girls and young women (or their friends) that lead them to suspect that individuals are at risk. Pressure may come from people other than the adult family members; it may be other children in the family who are pressurising one specific girl or young woman to undergo FGM.

Behavioural changes may indicate a risk of harm, or that harm has already occurred. A simple change such as prolonged visits to the toilet may indicate that a child is experiencing difficulties urinating following FGM.

This includes being aware of the possible true purpose of a girl's visit to the family's country of origin. Older girls and siblings may also be very aware of the risk or purpose of a planned visit abroad but be unable to protect themselves unaided. They may have confided in the practitioner, who must know how to operate within the safeguarding and legislative procedures.

While it is not the responsibility of individual practitioners to undertake investigations, they should be alert to considering FGM among general assessments and know who to refer concerns to. The concept of making every contact count (PHE, 2018) has been expanded in recent years, and FGM is one of the issues that should be considered. Where suspicion has been established a risk assessment needs to take place, be actioned and then documented. Figure 7 provides questions which may help with the risk assessment process, although each local area should have such systems in place that are understood by all frontline personnel who are likely to come in contact with women or girls affected by FGM.

Figure 7 – Questions to consider when undertaking a risk assessment

- Have I asked about FGM?
- If yes, what local services/support is available to meet her needs?
- Where does the woman/girl come from originally?
- Is the patient part of a community that practises FGM?
- Has she ever been cut or had any form of surgery or piercings?
- Is the girl/woman a victim/survivor of FGM?
- Is the girl at risk of FGM?
- Does she have daughter(s) under 18 years of age?
- Are there girls under 18 years of age at home?
- In discussing attitudes to FGM, do you conclude that it is more likely or less likely that she will subject her daughter(s) to FGM?
- Are there any plans to travel to a country where FGM is prevalent with girls in her family?
- Does she have sisters/other female relatives who have undergone FGM?
- Have I documented my concerns?
- What information do I need to share?

The Department of Health and Social care has produced a risk assessment process which can be found in its *Female Genital Mutilation Risk and Safeguarding Guidance for Professionals* (DH, 2016) at assets.publishing.service.gov.uk/media/5a8041f3ed915d74e622d655/FGM_safeguarding_report_A.pdf

Community, practice and travel nurses

Community and practice nurses, who have access to women in the community or home setting may note information leading them to think that girls may be at risk, such as behavioural changes. It is equally important for nurses working in travel, clinics or who come into contact with women who are travelling abroad to be vigilant, for example around passport authorisation.

Travel nurses need to understand that they may be in a position to identify girls who are at imminent risk of FGM, however this can be a challenging and complex area for travel health professionals because they may not have easy access to patient notes or the wider services that are available in acute trusts or GP surgeries. Travel health also encompasses many disciplines including nurses, doctors and pharmacists and is delivered in multiple settings in both NHS and the independent sector.

Nurses, nursing associates and midwives undertaking cervical screening tests should routinely assess patients for risk of FGM as part of the procedure. The experience of FGM

can access as a barrier to accessing screening and may make the screen test painful or not possible. Survivors of FGM should be offered a double appointment so that time is given to carry out the procedure with great care allowing for the support the patient may need. It may be necessary to refer the patient to a specialist clinic for the screen test (Jo's Trust, 2018).

Acute sector nurses

Nurses working across the full spectrum of acute services such as neonatal and child health, sexual health, accident and emergency, gynaecology, or other related areas should be aware of FGM. It is important to be able to respond appropriately in the best interests of anyone who may be at risk of abuse, or who may have already been mutilated.

In recognition of some of the specific issues faced in sexual health care the RCN published the following publications:

- *Female Genital Mutilation: RCN Guidance for Travel Health Services*
- *Female Genital Mutilation: RCN Guidance for Sexual Health Care*

These are available at: rcn.org.uk/clinical-topics/female-genital-mutilation/publications.

Midwives

Midwives are most likely to encounter women who have been mutilated, and it is important to ask the question during pregnancy to ensure a safe birth and postnatal care for both mother and baby. Maternity services, especially where there are known FGM practising communities, will have specialist midwives who take the lead on supporting these women and their colleagues in better understanding the issues surrounding FGM.

Midwives may also be concerned where baby girls are born to women who have had FGM performed, and this will naturally require a sensitive approach. There are differing views at present on whether this puts the girl at risk, but child protection must remain paramount. Documentation of conversations and concerns are vital to ensure better continuity of care going forward. However, if there are any concerns then further action is essential and can be carried out via safeguarding leads. Equally, midwives may also become concerned about a girl being at risk while attending a family for the birth of a subsequent child.

Figure 8 – Identifying girls at risk

This is difficult because:

- FGM happens only once
- parents may believe FGM is a good thing to do for their daughters
- the genitalia of girls are rarely examined
- it is not culturally acceptable for girls to talk openly about FGM.

There is a risk if:

- the girl's mother or her older sisters have been cut
- the woman or girl has limited contact with people outside of her family
- the paternal grandmother is very influential within the family
- the woman has poor access to information about FGM
- no one talks to the mother about FGM
- health, social service and education staff fail to respond appropriately
- communities are given the impression that FGM is not taken seriously by the statutory sector
- the woman has strong links with country of origin, family remaining in country of origin/frequent travel to country of origin.

Adapted from the Foundation for Women's Health, Research and Development (FORWARD) training pack, 2006

Registered nursing associates

Nursing associates on the NMC register have the same responsibilities as nurses and midwives in respect to FGM, as they have to adhere to the NMC Code (2018).

In the *Nursing Associate Standards* (2018) the following are noted:-

1.2 Understand and apply relevant legal, regulatory requirements, policies and ethical frameworks, including any mandatory reporting duties to all areas of practice.

3.24 Taken personal responsibility to ensure that relevant information is shared according to local policy and appropriate and immediate action is taken to provide adequate safeguarding and the concerns are escalated.

Confidentiality

To safeguard children and young people it may be necessary to give information to people working in other parts of the health service or outside of it. For some practitioners this can pose dilemmas, but both the law and policy allow for disclosure where it is in the public interest or where a criminal act has been perpetrated or a child is at risk. Parents

are responsible for their children and they may fear having this responsibility (or even the child) taken away from them. There may also be a perception that passing on information can damage the relationship of trust built up with families and communities. Nonetheless, it is crucial that the focus is kept on the best interests of the child as required by law. *The NMC's Code* (2018), clearly identifies that: 17: Raise concerns immediately if you believe a person is vulnerable or at risk and needs extra support and protection.

The NMC is also clear about professional responsibility, in particular around confidentiality. It is normally expected that information is shared with others only with the consent of the patient or client, but makes provision for when this is not possible 'if the patient or client withholds consent, or if consent cannot be obtained for whatever reason.

Disclosures may be made only where these:

- can be justified in the public interest (usually where disclosure is essential to protect the patient or client or someone else from the risk of significant harm)
- are required by law or an order of court.

The NMC Code (2018)

Respect people's right to privacy and confidentiality. As a nurse, midwife or nursing associate, you owe a duty of confidentiality to all those who are receiving care. This includes making sure that they are informed about their care and that information about them is shared appropriately.

To achieve this, you must:

5.1 respect a person's right to privacy in all aspects of their care

5.2 make sure that people are informed about how and why information is used and shared by those who will be providing care

5.3 respect that a person's right to privacy and confidentiality continues after they have died

5.4 share necessary information with other health and care professionals and agencies only when the interests of patient safety and public protection override the need for confidentiality

5.5 share with people, their families and their carers, as far as the law allows, the information they want or need to know about their health, care and ongoing treatment sensitively and in a way they can understand (NMC, 2018).

Referral to appropriate agencies

Caring for girls and women affected by FGM is not the primary responsibility of one professional group; it requires multi-agency management and nurses and midwives need to be aware of local procedures for referral where safeguarding is a concern. It is also important to understand local structures and services which may benefit a woman or

girl who has physical, psychological or psycho-sexual complications as a result of being mutilated.

Knowledge of local support groups or campaigns can be invaluable additions to the toolkit required to support best practice.

Education, raising awareness and conscious engagement to change

Raising awareness and consciousness about the practical, socio-cultural, ethico-legal, sexual health and practice care implications involved in FGM is essential, if real change is to happen.

It has long been recognised that the key to eliminating FGM is a cross-organisational, multiprofessional approach bringing together the expertise of survivors, activists, communities, the Police, social workers, teachers and health care professionals.

There is now increasing focus on engaging men and boys to exert their influence in families and communities as fathers and future husbands and decision makers and the role of men and boys in ending FGM was the focus for the 2023 International Zero day. In its report *Engaging Boys and Men to End FGM* (Unicef, 2023) outlines a range of multi-faceted projects to support male advocacy and presents data on changing societal attitudes, identifying that girls are less at risk of FGM if both parents oppose it.

Education and training needs to be provided for all health and social care professionals who may work with affected women and girls and with their families. It is also important to consider the issues of ethnicity, custom, culture and religion in a sensitive manner. Professionals should explore ways of resolving problems about the continuation of this practice in ways that involve women (and their communities) with their full participation.

The RCN believes that FGM should be a part of health education in all pre-registration and post-registration programmes for nurses, midwives, nursing associates and health visitors. It is also important for all registrants to ensure they are adequately prepared to provide effective practice in respect of FGM, in line with the NMC code (NMC, 2018). It is equally essential to raise awareness and the seriousness of the issues among teachers, school nurses and social service staff, as well as registered nurses, nursing associates and midwives. There are increasing opportunities locally and nationally for access to appropriate education, dependant on the expected level of engagement in the subject by health professionals.

The Department of Health and Social care commissioned Health Education England to produce e-learning training sessions. These are provided on the national eLearning for Health platform, available free to all NHS trusts (available at: e-lfh.org.uk/programmes/femalegenital-mutilation) and includes an introductory module for all health care staff.

As FGM is a safeguarding issue, it should be integral to all safeguarding training and annual updates to ensure all health care personnel have contemporary knowledge and a good understanding of identifying and referring any cases they come in contact with.

Practice matters and procedures

Professional curiosity and assessing need

Nurses and midwives need to be aware of how to sensitively care for women and girls, as well as being able to safeguard those at risk. They also need to be aware that accepting and respectful attitudes are vitally important to girls and women who have been cut. Equally, health professionals need to be curious about their local community to understand their needs and to actively engage with women to fully understand the community perspective on challenging issues such as FGM. Continuity of care and a holistic attitude to care provision will all support best practice.

The CMO's report (Davies, 2015) recommends that “all under-18s with suspected or confirmed FGM should be seen by or with a paediatrician with relevant expertise and experience, as well as knowledge of safeguarding”.

Women and girls who have suffered mutilation may be very reluctant to agree to a vaginal or rectal examination, and may refuse routine cervical smears and/or infection screening. It may be impossible to perform a vaginal examination at all, and can be very difficult or impossible to pass a urinary catheter. Nurses and midwives need to be compassionate and caring, exhibiting a sensitive manner, and be prepared sufficiently so that they do not exhibit signs of shock, confusion, horror or revulsion on seeing the genitalia.

Making Every Contact Count (PHE, 2018) is a programme focused on improving the lives of all those who come in contact with health care professionals, and is an ideal opportunity to develop a relationship with women and their families and to gather information about issues such as FGM. It is intended to create opportunities within existing health consultations to make every contact with a health professional count towards increasing health and well being awareness. Further information on Making Every Contact Count, public health topics and nursing roles is available from the RCN website at: rcn.org.uk

Despite the need for sensitivity, it is important to ask every woman whether they have been cut or circumcised. Some may seek help because they wish to have the FGM reversed before marrying, or may be experiencing problems conceiving because of difficulties with penetration and need to be referred to appropriate specialist clinics.

A deinfibulation service should be available, should be well-advertised via women's groups, and be easily accessible to those who may need it. Deinfibulation involves opening the scar tissue that covers the vaginal introitus and the urethral meatus surgically. Although best performed when not pregnant, women may need deinfibulation to be done as an emergency, for example, during a miscarriage. This is because products of conception, such as blood clots and fetal tissue, can be retained behind scar tissue and could lead to serious infection. It may also be performed during pregnancy in the first trimester or at time of birth to facilitate a safer vaginal birth.

Sometimes the term “reversal” is used but the effects of FGM can not be undone. The term deinfibulation that refers to the medical procedure to open the introitus is more accurate.

Knowledge about diversity

In our multi-cultural and multi-ethnic society, self-awareness and knowledge of diversity are essential skills for all health care professionals, enabling them to provide high quality individualised care. Concerns frequently articulated when giving reasons for why it may be difficult to engage with girls and women who need safeguarding because of FGM is not wishing to appear to act in discriminatory ways or from racist motivations.

FGM is mostly found in Africa, the Middle East and Asia, and while cultural sensitivity to the girl, woman and her family is always paramount, it should not override the safety or wellbeing of individuals. The inquiry report into the death of Victoria Climbié clearly notes the danger of making assumptions about cultural background that conflict with ensuring children's safety (House of Commons Health Committee, 2003). Lord Laming noted that children's needs for protection are the same whatever their cultural background, saying:

“a child is a child regardless of colour – if we are not careful we'll lose the whole emphasis on the child's welfare.”

(House of Commons Health Committee, 2003)

There is growing recognition of the need to better understand our gender diverse society that includes people who identify as trans-gender, non-binary or gender fluid. The umbrella term “trans” relates to people who have a gender identity that is not congruent with the gender that they were assigned at birth, and those who are non-binary have a gender identity that does not fit the binary male/female definition. It is important that health care professionals recognise gender diversity and ensure that their communication is sensitive to this. People who do not identify as female may have been affected by FGM and careful assessment should be undertaken to ensure that the needs of all survivors are met regardless of how they identify.

Antenatal care and reversal of infibulation (deinfibulation)

It is important to identify women who have been cut when they first seek pregnancy care, and find out what type of FGM has been performed. It will be necessary to ask about FGM as they may not volunteer the information.

Apart from the usual screening and antenatal care, it is important to provide pregnant women with support specific to their needs around FGM. They may need counselling, advice, information and social and psychological support. It is equally important to remember that a pregnant woman under the age of 18 will be subject to the mandatory duty to report and safeguarding processes if found to have been subjected to FGM. Teenagers may need extra support, particularly as there will be safeguarding issues to be addressed.

It is incumbent for health care professionals to complete documentation in relation to mandatory reporting and information sharing to protect other family members who may be at risk and to enable future service planning.

Deinfibulation (which is sometimes referred to as reversal, however the damage caused by FGM cannot be reversed) is best performed before or at least within the second

trimester of pregnancy at around 20 weeks of gestation. This avoids the need to cut the scar tissue in labour and reduces the possibility of extensive lacerations that can occur when the fetal head stretches the scarred or closed introitus and perineum. These may involve the urethra, bladder and rectum if uncontrolled and leave the woman with a fistula. It will also reduce the risk of fetal asphyxia or stillbirth if a woman progresses unaided to the second stage of labour.

Surgical deinfibulation should be offered where appropriate. Partners should be involved in decision making when the woman is willing for this to happen. It is important to work out an agreed care plan with the woman early in pregnancy, and involve trained professional interpreters to ensure that medical information is correctly presented. Caesarean section is not indicated just because a woman has had FGM performed. The midwife should always assess the need for an elective episiotomy in labour.

It must be noted, however, that women may be reluctant to undergo deinfibulation until labour commences, because this may be normal practice in their country of origin. This reinforces the importance of careful and sensitive explanation in pregnancy of why antenatal deinfibulation is preferable. It also underlines the importance of all midwives understanding what to do in this situation. It is also essential to inform women that they may still need a standard (posterior) medio-lateral episiotomy for fetal distress in the second stage of labour. This should be explained because women may be very disappointed if they have to have perineal suturing after the birth, despite having had a deinfibulation in pregnancy.

The aim of deinfibulation is to restore normal anatomy as far as possible, which may be very limited depending on the damage caused originally. The procedure is the same in principle whether it is carried out as an elective procedure before pregnancy, in the antenatal period, or in labour itself. It can be performed by a midwife if necessary during the first stage of labour once the presenting part is low (RCOG, 2015).


Principles of good practice include:

- adequate pain relief (general, regional or local anaesthesia) is essential; non pregnant women may prefer to have general anaesthesia because the procedure can bring back very traumatic memories of when they were cut, however, this may not be recommended or possible when pregnant or in labour
- using aseptic techniques following cleansing of the vulval area; also pay careful attention to hand washing and wear sterile gloves
- examine the vulval area carefully, infiltrate with local anaesthetic and then open the scar in the midline, exposing the underlying tissues which may include the clitoris
- a midline incision along the scar is less likely to bleed heavily and will follow a line that may already have areas of weakness where the original healing of the edges was incomplete; it can be easier to do this if the tissue is carefully lifted along the midline with a finger or blunt instrument
- if the clitoris is present and can be palpated, an experienced practitioner can extend the incision to expose the clitoris and free any para-clitoral adhesions; if uncertain, cutting should stop when the urinary meatus is visible

- suture the raw edge on each side of the labia with fine dissolvable sutures to ensure haemostasis and an over-sewing stitch; this is important also to ensure the raw edges do not fuse together
- provide adequate analgesia following the deinfibulation
- if there is extensive fibrosis of the vaginal introitus, perhaps from the use of corrosive substances or angurya cuts, an episiotomy may be needed
- provide advice on good personal hygiene, especially keeping the area clean couples should be advised to avoid intercourse until healing has occurred and to use a lubricant if necessary
- women need to be advised that urine and menstrual flow may appear heavier because of the removal of the scar-tissue barrier.

The deinfibulation procedure is illustrated in Diagram 7.

Diagram 7 – Cutting open the scar



Care in labour

Principles of good practice include:

- normal care is required during the first stage of labour; usual sensitivity is essential
- there is no need to pass a catheter unless the woman is unable to pass urine
- deinfibulation may need to be carried out during the first stage of labour or at the time of delivery
- midwives need to watch women who have undergone type 3 FGM closely during the second stage of labour, even when the woman's introitus has previously been assessed as adequate for the birth; unexpected problems may occur with descent of the fetal head or stretching of the perineum because the scar tissue around the vagina and perineum may be unstable

- a medio-lateral episiotomy should be performed in the second stage of labour only if unavoidable
- it is important to explain the requirements of UK law
- it is not acceptable to reinfibulate or stitch the woman back closed after the birth.

If reversal of the infibulation has not already been performed, it needs to be carried out during the first stage of labour using adequate analgesia. If the second stage has already been reached, a midline incision must be used.

Suturing a laceration or an episiotomy It is of course important to repair a laceration or episiotomy, to stem the bleeding. If the woman did not have a deinfibulation during the antenatal period, midwives often perform an anterior episiotomy and this will require careful repair. The key is to ensure that both the urethra and introitus are able to be seen following repair.

Re-suturing or reinfibulation or closing should never be considered or offered. This may mean that careful discussions have to be held with the woman and family to explain the law and why reinfibulation has to be refused. Women may themselves request reinfibulation for social reasons or because they have known nothing else.

It is necessary to follow up with the woman during the postnatal period. Support, information and counselling continue to be very important.

Health care professionals who participate in FGM or reinfibulation risk both criminal charges and fitness to practise proceedings before the NMC.

The most important points to remember are to:

- ensure early identification – during antenatal booking or first visit
- arrange for deinfibulation during the first stage of labour with adequate pain relief
- support the woman with sensitivity
- notify her health visitor, GP and other professionals if the baby born is a girl, with regard to safeguarding the child; this is also to provide ongoing information and support to the family
- continue to provide postnatal support
- consider referring to an organisation that can offer additional support and information.

Conclusion

FGM is a global and national issue, although significant progress has been achieved, there are many challenges ahead that must not be underestimated. In many cities across the UK there are pockets of good practice, support groups and professional expertise which is well established and accessible. Nevertheless, there is a continuing need to extend services and provide resources to accommodate the needs of women, girls, families and communities to support their health and wellbeing.

Education is a critical tool in the fight to change attitudes to FGM and eradicate this violence from society; teaching about FGM and its consequences should be imbedded in school education, as well as pre- and post-registration education for all those involved in caring for girls and women in local communities.

Women and girls who have been abused need particular and sensitive support, together with access to facilities to help them with the physical, psychological and social consequences of this potentially devastating abuse. All professionals, practising communities and the public have a role to play in this social change towards eradication. Change can only take place to keep women and girls safe if practising communities are involved at all stages of child protection and service provision.

FGM also demands professional curiosity and awareness by those who come in contact with girls and women, including those health care professionals who may themselves have come from FGM practising communities. A further challenge for many is to consider how awareness raising can be enhanced simply by voicing concerns, speaking to colleagues and engaging in the campaign to end this violation of basic human rights. This should also include being sensitive to how students or colleagues may react to learn that such mutilation takes place. Nurses and midwives have a role to play by being politically engaged in movements that impact on care, and this is particularly so with FGM.

FGM is a complex issue and requires vigilance, as well as continuing engagement with the legislation, and the changing scene of caring for girls and women who may be at risk or may have been mutilated by FGM.

Tackling FGM requires a multi-agency and multiprofessional approach and response through a recognised pathway that supports quality, evidence-based care and safeguarding. It is the responsibility of all nurses and midwives, working both in city and rural locations, to recognise FGM as abuse and to know who their local contacts are for support, training and action should they have a concern.

Highlighting the campaign to end FGM

The campaign to end FGM remains a constant challenge and nurses, midwives and registered nursing associates have much to contribute to this by remaining vigilant to all harmful practices including so called honour-based abuse, forced marriage, virginity testing, hymenoplasty and FGM which violate the basic human rights of women and girls. Health care professionals have a responsibility to keep up to date so that they are equipped to spot signs, initiate courageous and sensitive conversations, to safeguard and to ensure survivors have the specialist care that they need.

References and further reading

African Union (2003) *Protocol to the African charter on human and people's rights on the rights of women in Africa*, Addis Ababa: AU. Available at: www.un.org/shestandsforpeace/content/protocol-african-charter-human-and-peoples-rights-rights-women-africa (accessed 5 October 2023).

Bagness C (2015) Tackling FGM – everybody's business, *Practice Nursing*. Published Online: 31 Jan 2015. Available at: <https://doi.org/10.12968/pnur.2015.26.2.74> (accessed 5 October 2023).

Bagness C (2019) FGM: raising awareness in community-based health practice, *Practice Nursing*. Available at: www.magonlinelibrary.com/doi/abs/10.12968/pnur.2019.30.10.502 (accessed 5 October 2023)

Brown E and Hemmings J (2013) *The FGM initiative: evaluation of the first phase (2010-2013)*, Options UK: London. Available at: www.unicef.org/media/106831/file/FGM-State-of-Evidence.pdf (accessed 5 October 2023).

Chiodini JH, Anderson E, Driver C, Field VK, Flaherty GT, Grieve AM, Green AD, Jones ME, Marra FJ, McDonald AC, Riley SF, Simons H, Smith CC and Chiodini PL (2012) Recommendations for the practice of travel medicine, *Travel Medicine and Infectious Disease*, 10 (3), pp.109-28. Available at: pubmed.ncbi.nlm.nih.gov/22658381 (accessed 5 October 2023).

Davies SC (2015) *Annual Report of the Chief Medical Officer, 2014, The Health of the 51%: Women*, London: DHSE. Available at: assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/595439/CMO_annual_report_2014.pdf (accessed 5 October 2023).

DHSE (2014) *New FGM measures launched to 'care, protect, prevent'* (part of Girl Summit 2014: Improving the lives of girls and women in the world's poorest countries), London: DHSE. Available at: www.gov.uk/government/news (accessed 5 October 2023).

DHSE (2015) *Female Genital Mutilation Risk and Safeguarding: Guidance for professionals*. Available at: assets.publishing.service.gov.uk/media/5a8041f3ed915d74e622d655/FGM_safeguarding_report_A.pdf (accessed 5 October 2023).

DHSE (2015) *Female genital mutilation (FGM): guidance for healthcare staff*. Available at: www.gov.uk/government/collections/female-genital-mutilation-fgm-guidance-for-healthcare-staff (accessed 5 October 2023).

DHSE (2015) *FGM enhanced dataset: guidance on NHS staff responsibilities*. Available at: www.gov.uk/government/publications/fgm-enhanced-dataset-guidance-on-nhs-staff-responsibilities (accessed 5 October 2023).

DHSE (2015) *Commissioning services for women and girls with FGM*. Available at: www.gov.uk/government/publications/services-for-women-and-girls-with-fgm (accessed 5 October 2023).

DHSE (2017) *Safeguarding women and girls at risk of FGM*. Available at: www.gov.uk/government/publications/safeguarding-women-and-girls-at-risk-of-fgm (accessed 5 October 2023).

DHSE (2022) *Virginity testing and hymenoplasty: multi agency guidance*. Available at: www.gov.uk/government/publications/virginity-testing-and-hymenoplasty-multi-agency-guidance/virginity-testing-and-hymenoplasty-multi-agency-guidance (accessed 5 October 2023).

El Dirani Z, Farouki L, Aki C, Ali U, Akik C and McCall S (2022) Factors Associated with FGM: a systematic review and synthesis of national regional and community based studies. *BMJ Sexual and Reproductive Health Journal*. Jul 2022 48 (3) 169-178. (accessed 5 October 2023).

END FGM European Campaign (2009) *Ending female genital mutilation: a strategy for the European Union institutions*, Dublin: Amnesty International. Available at: www.endfgm.eu (accessed 5 October 2023).

FORWARD (2023) *Female genital mutilation as grounds for asylum*. Available at: www.forwarduk.org.uk (accessed 5 October 2023).

Gov.UK (2015) *What to do if you're worried a child is being abused, Advice for practitioners*. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/419604/What_to_do_if_you_re_worried_a_child_is_being_abused.pdf (accessed 5 October 2023).

Gov.UK (2016) *Multi-agency statutory guidance on female genital mutilation*, London: FCO. Available at: www.gov.uk/government/publications/multi-agency-statutory-guidance-on-female-genital-mutilation (accessed 5 October 2023).

Gov.UK. Children's services safeguarding. Available at: www.gov.uk/childrensservices/safeguarding-children (accessed 5 October 2023).

Gov.UK (2014) *There is no justification for FGM – it is child abuse and it is illegal* (Government declaration to end FGM in the UK and abroad, issued to mark the International Day of Zero Tolerance to Female Genital Mutilation, 6 February 2014). Available at: www.gov.uk/government/news/new-government-measures-to-end-fgm (accessed 5 October 2023).

Gov.UK (2021) *Tackling violence against women and girls*. Available at: www.gov.uk/government/publications/tackling-violence-against-women-and-girls-strategy (accessed 5 October 2023).

Gov.UK (2022) *Working Together to Safeguard Children*. Available at: www.gov.uk/government/publications/working-together-to-safeguard-children--2 (accessed 5 October 2023).

Gov.UK (2023) *Female Genital Mutilation Act 2023*. Available at: www.legislation.gov.uk/ukpga/2003/31 (accessed 5 October 2023).

Gov.UK (2015) *Mandatory reporting of female genital mutilation: procedural information*. Available at: <https://www.gov.uk/government/publications/mandatory-reporting-of-female-genital-mutilation-procedural-information> (accessed 5 October 2023).

Home Office (2014a) *Ending violence against women and girls in the UK* (Policy), London: Home Office. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/229477/VAWG_Brief_v3.pdf (accessed 5 October 2023).

House of Commons Health Committee (2003) *The Victoria Climbié inquiry report: sixth report of session 2002-3, report and formal minutes together with oral evidence*, London: TSO. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/273183/5730.pdf (accessed 5 October 2023).

Hussein E (2010) *Women's experiences, perceptions and attitudes to female genital mutilation: the Bristol PEER study*, London: Forward. Available at: www.forwarduk.org.uk/forward-publications/womens-experiences-perceptions-and-attitudes-of-female-genital-mutilation (accessed 5 October 2023).

Information Standards Board for Health and Social Care (2014) ISB 1610 female genital mutilation prevalence dataset: standard specification, ISB: Leeds. Available at: <https://digital.nhs.uk/data-and-information/information-standards/information-standards-and-data-collections-including-extractions/publications-and-notifications/standards-and-collections/scci2026-female-genital-mutilation-enhanced-dataset> (accessed 5 October 2023).

Jo's Trust (2023) *Let's talk about cervical screening and FGM*. Available at: www.jostrust.org.uk/about-us/news-and-blog/blog/smear-tests-fgm (accessed 5 October 2023).

Liao LM, Elliott C and Creighton SM (2013) Adult recall of childhood female genital cutting and perceptions of its affects: a pilot study for service improvement and research feasibility, *Journal of Obstetrics and Gynaecology*, 33(3), pp.292-295. Available at: <https://pubmed.ncbi.nlm.nih.gov/23550862> (accessed 5 October 2023).

Macfarlane and Dorkenoo (2014) *Female genital mutilation in England and Wales: updated statistical estimates of the numbers of affected women living in England and Wales and girls at risk*. Interim report on provisional estimates, London: City University London. Available at: <https://openaccess.city.ac.uk/id/eprint/3865> (accessed 5 October 2023).

Muslim Council of Britain (2014) Muslim Council of Britain speaks out against female genital mutilation (Statement and news release, 23 June 2014). Available at: www.mcb.org.uk (accessed 5 October 2023).

Northern Ireland Department of Health, Social Services and Public Safety (2014) *Multi-agency practice guidelines: female genital mutilation*, Belfast: DHSSPSNI. Available at: www.health-ni.gov.uk/publications/multi-agency-practice-guidelines-female-genital-mutilation-0 (accessed 5 October 2023).

Nursing and Midwifery Council (2018) *The Code: Professional standards of practice and behaviour for nurses, midwives and nursing associates*, London: NMC. Available at: www.nmc.org.uk/standards/code/read-the-codeonline (accessed 5 October 2023).

Nursing and Midwifery Council (2022) *Safeguarding hub*. Available at: www.nmc.org.uk/standards/guidance/safeguarding (accessed 5 October 2023).

Nursing and Midwifery Council (2018) *Standards for nursing associates*, NMC: London. Available at: www.nmc.org.uk/standards/standardsfor-nursing-associates (accessed 5 October 2023).

National Society for the Prevention of Cruelty to Children. *Female genital mutilation factsheet*, London: NSPCC. Available at: www.nspcc.org.uk (accessed 5 October 2023).

Gov.UK (2015) *Serious Crime Act*. Available at: www.gov.uk/government/collections/serious-crime-bill (accessed 5 October 2023)

Public Health England (2018) *Making Every Contact Count*. Available at: www.makingeverycontactcount.co.uk (accessed 5 October 2023).

Royal College of Midwives, Royal College of Nursing, Royal College of Obstetricians & Gynaecologists, Equality Now, UNITE (2013) *Tackling FGM in the UK: intercollegiate recommendations for identifying, recording, and reporting*, London: RCM. Available at: www.rcn.org.uk/Professional-Development/publications/pub-004531 (accessed 5 October 2023).

Royal College of Nursing (2021) *Safeguarding children and young people – every nurse's responsibility*, London: RCN. Available at: www.rcn.org.uk/Professional-Development/publications/safeguarding-children-and-young-people-every-nurses-responsibility-uk-pub-009-507 (accessed 5 October 2023).

Royal College of Paediatrics and Child Health (2019) *Safeguarding children and young people: roles and competences for health care staff. Intercollegiate document* (Third edition), London: RCPCH. Available at: www.rcpch.ac.uk/resources/safeguarding-children-young-people-roles-competencies (accessed 5 October 2023).

Royal College of Obstetricians and Gynaecologists (2015) *Female genital mutilation and its management* (Green-top guideline No.53), London: RCOG. Available at: www.rcog.org.uk/guidance/browse-all-guidance/green-top-guidelines/female-genital-mutilation-and-its-management-green-top-guideline-no-53 (accessed 5 October 2023).

Scottish Government (2005) *Prohibition of Female Genital Mutilation (Scotland) Act 2005* (2005 asp8), Edinburgh: TSO). Available at: www.legislation.gov.uk/asp/2005/8/contents (accessed 5 October 2023).

Scottish Government (2014a) *Violence against Women and Girls (VAWG)*. Available at: www.gov.scot/policies/violence-against-women-and-girls (accessed 5 October 2023).

United Nations (1948) *The universal declaration of human rights*, New York: UN. Available at: www.un.org/en/about-us/universal-declaration-of-human-rights (accessed 5 October 2023).

United National Population Fund and United Nations Children's Fund (2013) *UNFPA-UNICEF joint programme on female genital mutilation/cutting: annual report 2012*, New York: UNFPA and UNICEF. Available at: www.unfpa.org/publications/unfpa-unicef-joint-programme-female-genital-mutilationcutting-annual-report-2012 (accessed 5 October 2023).

UNICEF (2023) *FGM Facts*. Available at: <https://data.unicef.org/topic/child-protection/female-genital-mutilation> (accessed 5 October 2023).

UNICEF (2023) *Engaging boys and men to end female genital mutilation* (web). Available at: <https://data.unicef.org/resources/mobilizing-boys-and-men-in-the-fight-against-female-genital-mutilation> (accessed 5 October 2023).

World Health Organization (2001) *Female genital mutilation: the prevention and the management of the health complications. Policy guidelines for nurses and midwives*, Geneva: WHO. Available at: www.who.int/publications/i/item/WHO-FCH-GWH-01.5 (accessed 5 October 2023).

World Health Organization (2010) *Global strategy to stop health-care providers from performing female genital mutilation* (UNAIDS, UNDP, UNFPA, UNHCR, UNICEF, UNIFEM, WHO, FIGO, ICN, MWIA, WCPTA, WMA), Geneva: WHO. Available at: www.who.int/publications/i/item/WHO-RHR-10.9 (accessed 5 October 2023).

World Health Organization (2012) *Understanding and addressing violence against women: female genital mutilation* [Information sheet], Geneva: WHO. Available at: www.who.int (accessed 5 October 2023).

World Health Organization (2018) *Female genital mutilation: fact sheet*, Geneva: WHO. Available at: www.who.int/en/news-room/fact-sheets/detail/female-genital-mutilation (accessed 5 October 2023).

World Health Organization (2023) *FGM Factsheet*. Available at: www.who.int/news-room/fact-sheets/detail/female-genital-mutilation (accessed 5 October 2023).

Resources

Things not to say to FGM campaigners and survivors. [youtube.com/watch?v=BJES6Usn8eQ](https://www.youtube.com/watch?v=BJES6Usn8eQ)

Daughters of Eve. lighthousevictimcare.org/organisation/daughters-of-eve

#EndFGM. It's our time now. [youtube.com](https://www.youtube.com)

#EndFGM. Our daughters. [youtube.com](https://www.youtube.com)

#EndFGM. The Words Don't Come. [youtube.com](https://www.youtube.com)

Forward UK. forwarduk.org.uk

The Girl Generation. thegirlgeneration.org

Health Education England. FGM e-learning module. e-lfh.org.uk

Health Education England. Cultural competence e-learning module. e-lfh.org.uk

UK Government resources can be found at: gov.uk

Let's Talk FGM App. oxfordhealth.nhs.uk/support-advice/getting-help/lets-talk-fgmapp

National Society for the Prevention of Cruelty to Children (NSPCC) 0800 028 3550 or fgmhelp@nspcc.org.uk. nspcc.org.uk/what-is-child-abuse/types-of-abuse/female-genital-mutilation-fgm

NHS England resources on FGM can be found at: nhs.uk/conditions/female-genital-mutilation-fgm

NHS England. Six videos available. [youtube.com](https://www.youtube.com)

Petals. App containing information about FGM, personal stories from those who have been affected, links to educational films, a quiz and tips on how to get involved in campaigning to end the practice. <http://petals.coventry.ac.uk>

Petals for Professionals. App provides access to information and knowledge about FGM; the health impacts; the legal responsibilities of professionals; advice on initiating conversations; information for specific professions; and where to get support and advice. <http://petals.coventry.ac.uk/professionals>

Royal College of Nursing. Female genital mutilation. rcn.org.uk/clinical-topics/female-genital-mutilation

Royal College of Obstetricians and Gynaecologists resources on FGM can be found at: rcog.org.uk/for-the-public/browse-our-patient-information/female-genital-mutilation-fgm

Royal College of Paediatrics and Child Health resources on FGM can be found at: rcpch.ac.uk/resources/female-genital-mutilation-resources

Safeguarding Board for Northern Ireland. safeguardingni.org/search?text=FGM

Virtual College. A free online FGM safeguarding training course developed for the Home Office by Virtual College. virtual-college.co.uk/resources/free-courses/recognising-and-preventing-fgm

World Health Organization (2023) *FGM Factsheet*. who.int/news-room/fact-sheets/detail/female-genital-mutilation

Organisations and support groups

England

Black Women's Health and Family Support (BWHAFS). nhs.uk/services/service-directory/black-women-s-health-and-family-support-bwhafs/N10498418

London Safeguarding Children Procedures and Practice Guidance. londonsafeguardingchildrenprocedures.co.uk

Refugee Council. refugeecouncil.org.uk/about-us/contact

Northern Ireland

The Northern Ireland Council for Racial Equality (NICRE). <https://nicre.org/about-us/>

Scotland

Save the Children Scotland. savethechildren.org.uk/scotland

Scottish Refugee Council. scottishrefugeecouncil.org.uk

Wales

Bawso. bawso.org.uk

Welsh Refugee Council. <https://wrc.wales>


National and international groups

FGM National Group. fgmnationalgroup.org

FORWARD (Foundation for Women's Health, Research and Development).
forwarduk.org.uk

Appendices

Appendix 1: Map showing the global distribution of FGM



Appendix 2: Alternative terms used for FGM

Country	Term used for FGM	Language	Term used for FGM	Language
CHAD – the Ngama sara subgroup	Bagne		Gadja	
EGYPT	Thara, Khitan, Khifad	Arabic		
ETHIOPIA	Megrez	Amharic	Absum	Harrari
ERITREA	Mekhnishab	Tigreña		
GAMBIA	Niaka, Kuyango, Musoluta Karoota	Mandinka		
GUINEA-BISSAU	Fanadu di Mindjer	Kriolu		
IRAN	Xatna	Farsi		
KENYA	Kutairi ; Kutairi was ichana	Swahili		
NIGERIA	Ibi/Ugwu	Igbo	Didabe fun omobirin / ila kiko fun omobirin	Yoruba
SIERRA LEONE	Sunna	Soussou	Bondo	Temene
	Bondo/Sonde	Mende	Bondo	Mandinka
	Bondo	Limba		
SOMALIA	Gudinlin; Halalays; Qodin	Somali		
SUDAN	Khifad, Tahoor	Arabic		
TURKEY	Kadin Sunneti	Turkish		

Adapted from: HM Government (2018) Multi-agency statutory guidance on female genital mutilation.
[gov.uk/government/publications/multi-agency-statutory-guidance-on-female-genital-mutilation](https://www.gov.uk/government/publications/multi-agency-statutory-guidance-on-female-genital-mutilation)

Appendix 3: Department of Health NHS England Overview of the FGM mandatory reporting duty



Department
of Health


'Care, Protect, Prevent'

#EndFGM

NHS

England

FGM Mandatory reporting duty




If a girl appears to have been recently cut or you believe she is at imminent risk, act immediately – this may include phoning 999.

REMEMBER: Mandatory reporting is only one part of safeguarding against FGM and other abuse. **Always ask your local safeguarding lead if in doubt.**

Appendix 4: Tackling FGM in England

Diagrammatic representation of how government departments and NHS in England work together to ensure all nurses and midwives understand their role, and work collaboratively across different agencies to best support women and girls affected by FGM.



RCN quality assurance

Publication

This is an RCN practice guidance. Practice guidance are evidence-based consensus documents, used to guide decisions about appropriate care of an individual, family or population in a specific context.

Description

This comprehensive guidance considers many aspects of FGM, including contemporary socio-cultural, legal and clinical components to aid raising awareness and improving the quality of care to those who have been adversely affected by FGM.

Publication date: November 2023 Review date: February 2027

The Nine Quality Standards

This publication has met the nine quality standards of the quality framework for RCN professional publications. For more information, or to request further details on how the nine quality standards have been met in relation to this particular professional publication, please contact **publications.feedback@rcn.org.uk**

Evaluation

The authors would value any feedback you have about this publication. Please contact **publications.feedback@rcn.org.uk** clearly stating which publication you are commenting on.

The RCN represents nurses and nursing, promotes
excellence in practice and shapes health policies

RCN Direct
www.rcn.org.uk/direct
0345 772 6100

Published by the Royal College of Nursing
20 Cavendish Square
London
W1G 0RN

020 7409 3333
www.rcn.org.uk

November 2023
Publication code: 011 168

