Recovery experiences of patients under 60 with fragility hip fracture: A qualitative study

Dr Gillian Janes

Bradford Teaching Hospitals NHS Foundation Trust

Professor Laura Serrant

Sheffield Hallam University

Professor Magi Sque

University of Wolverhampton

Defining fragility hip fracture

Hip fracture: All fractures of the proximal femur (Archibald 2003)

Fragility fracture: A fracture resulting from lowenergy injury e.g. fall from standing height (Oetgen et al 2009; Chesser et al 2011)

Background

A serious injury, often due to a sudden, traumatic event, characterised by a complex recovery journey.

Relatively uncommon in young adults but:

- Significance is underestimated (Boden et al 1990)
- Potentially profound societal and individual social and economic implications (Holt et al. 2008)
- Can have devastating effects on patient and family (Young and Resnick 2009, Vilardo and Shah 2011)
- Minimally represented group in the literature, social discourse and healthcare policy, with little known of their recovery experiences and long-term outcomes

(Janes et al 2018)

Aims


- 1. To explore the recovery experiences of under 60s and identify potential service improvements
- 2. To test The Silences Framework

Methodology and Method


- Criticalist, anti-essentialist philosophy
- Interpretive methodology
- Guided by The Silences Framework:
 - Stage 1 Working in silences
 - Stage 2 Hearing silences
 - Stage 3 Voicing silences
 - Stage 4 Working with silences

Participant characteristics

Time since injury	No. participants
1-4yrs	14
5-9yrs	13
10yrs	3



Participant characteristics



Findings

One cross-cutting theme: *Communication*Four main themes:

Experience of care

Impact on self

Impact on others

Moving forward

Wide variation in care quality

'He was fantastic – he listened to me then did something' (AVS)

'Once discharged you're in no-man's land'

'...quite horrendous...you weren't nursed you were processed...worst time of my life...the lack of care really...lack of humanity' (RP) '...I don't fit the classic demographic for that kind of injury...so it might not be the first port of call for diagnosis...people that are under 50...' (GA)

Challenging social & professional norms

'A fit young man like you doesn't go around breaking your hip. (ST)

'...you don't present like a fractured hip...one of them looked at me and said have you always had a problem with falling Mrs...I think they actually thought I was...an alcoholic.' (SJP)

Loss (...and finding)

'got to stay where I am doing what I'm doing...it's a loss...that was my chance ...don't think it'll come again (DH)

'Car to bar [shoes]...it's like a loss of femininity for a woman.' (DM)

'Blessing in disguise...I've found myself again.' (GT)

'...they were...saying I was playing on it...I should be able to...walk...they said "I've had a hip replacement and it's a six-week recovery"...' (LL)

Post traumatic stress disorder symptoms

'I was quite an active person and...I have this fear of falling over...and that's sad...its very difficult to explain...what happens to me...I become very vulnerable...but I couldn't say to anybody I am absolutely terrified, petrified'. (MES)

'... such a lot of pain...I struggled...even now, sometimes it's so hard...I just keep going because I need my wages...feel like I've aged fifteen years...' (DH)

Resilience

'I just had to get on with it...nobody else could do it for me.' (RP)

'...I asked about physiotherapy...said I...can't do my job the way I am...he just dismissed it...said you're a motivated man you won't need it...there wasn't no plan...you just go do it kind of thing...' (MS)

Looking to the future

'...getting myself into a position where I've got support systems...financial outgoings are minimal...I've got something to fall back on...' (SF)

'...limited shelf-life' (GT)

'...didn't think it would affect me permanently...still have this half-hearted hope that one day...things will be as they were...think I'm deluding myself a bit really but don't want to lose that...' (VJJ)

In conclusion

- Improved public & professional awareness of fragility fracture in the under 60s is needed
- Long term multi-faceted impact was very common
- Listening to & involving patients & families is crucial
- Nurses must tailor care to the needs of younger patients
- Further research to address the gap in the literature, policy & professional discourse is needed

Thank you

Email: Gillian.Janes@bthft.nhs.uk Twitter: @DrGillianJanes

l.serrant@shu.ac.uk

@lauraserrant

m.sque@wlv.ac.uk

magi.sque@nhs.uk

Acknowledgements:

Study participants
Louise Donnison
Mr Ben Ollivere & Nottingham University Hospitals NHS Trust
School of Health & Social Care, Teesside University

References

- Archibald G. (2003) Patients' experiences of hip fracture. *Journal of Advanced Nursing*, **44(4**), 385-392
- Boden S.D., Labropoulos P. & Saunders R. (1990) Hip fractures in young patients: is this early osteoporosis? Calcified Tissue International 46(2), 65-72
- Chesser T.J.S., Handley R. & Swift, C. (2011) New NICE guideline to improve outcomes for hip fracture patients. *Injury* 42(8), 727-729
- Holt G., Smith R., Duncan K., Hutchison J.D.& Gregori A. (2008) Epidemiology and outcome after hip fracture in the under 65s evidence from the Scottish hip fracture audit. *Injury* 39(10),1175-1181
- Janes G (2016) Silent slips, trips and broken hips: The recovery experiences of young adults following an isolated fracture of the proximal femur. PhD Thesis. University of Wolverhampton
- Janes G, Serrant L, Sque M (2018) International Journal of Orthopaedic & Trauma Nursing https://doi.org/10.1016/j.ijotn.2018.02.006
- Janes G, Serrant L, Sque M Screaming Silences: Lessons learned from the application of a new research framework Nurse Researcher (in press)
- Oetgen M.E., Miki R.A., Smart L.R. & Lindskog D.M. (2009) Evaluation of bone mineral density and metabolic abnormalities associated with low-energy hip fractures. *Current Orthopaedic Practice* 20(6), 674-68
- Serrant-Green L (2010) The sound of 'silence': A framework for researching sensitive issues or marginalised perspectives in health *Journal of Research in Nursing* 16(4), 347-360
- Vilardo L & Shah M. (2011) Chronic pain after hip and knee replacement. <u>Techniques in Regional</u>
 <u>Anaesthesia and Pain Management</u> 15(3),110-115
- Young Y. & Resnick B. (2009) Don't worry, be positive: improving functional recovery 1 year after hip fracture. Rehabilitation Nursing 34(3),110-117